

高エネルギーガンマ線 観測の現状と将来

Status and future of high-energy Gamma-ray Astrophysics



森 正樹

東京大学宇宙線研究所

宇宙線研究所将来計画に向けた勉強会




2006年4月20日 宇宙線研究所

Detection of gamma-rays (1)




Atmospheric Cherenkov telescopes


Cherenkov light from gamma-ray showers
Lateral distribution \wp *Timing distribution*




Atmospheric transmission



MODRAN4 calculation by M.Yuasa (2006)




MODRAN4 calculation by H. Tsunoo (2002)





Imaging Cherenkov Telescopes

Shower profile




Focal plane image




α (image orientation angle)

Imaging analysis



D.J. Fegan, J.Phys.G, 1997

Distribution of imaging parameters




(Simulation)

α (image orientation angle)

"Image parameters": A.M. Hillas, 1983 ICRC

"Alpha": A.V.Plyasheshnikov and G.F.Bignami, N.C. 1985

WhippleによるCrabの検出




82hrs, 0.24 γ /min


Weekes et al. ApJ 342 (1989) 349

WhippleによるMrk421の検出

ガンマ
線信号



Stereo observation



© S.Funk, 2005


Angular resolution

$0.25\text{deg} \rightarrow 0.1 \text{ deg}$

Energy resolution

$30\% \rightarrow 15\%$

Better S/N (no local muons)



Detection of gamma-rays (2)

Base	Satellite	Ground	Ground
Gamma-ray detection	Direct (pair creation)	Indirect (atmospheric Cherenkov)	Indirect (shower array)
Energy	< 30 GeV \rightarrow 100 GeV	>100 GeV \rightarrow 50 GeV	>3 TeV \rightarrow 1 TeV
Pros	High S/N Large FOV	Large area Good $\Delta\theta$	24hr operation Large FOV
Cons	Small area High cost	Low S/N (CR bkgd.) <i>(but imaging overcomes this!)</i> Small FOV	Low S/N (CR bkgd.) Moderate $\Delta\theta$

VHE Experimental World

MILAGRO



STACEE



MAGIC



TIBET



MILAGRO

STACEE

VERITAS



MAGIC

TACTIC

TIBET
ARGO-YBJ

PACT

GRAPES

TACTIC



HESS



CANGAROO



Table 1. Currently operating VHE gamma-ray telescopes. The name of each telescope is given, along with its type (AC=Atmospheric Cherenkov), location, altitude, specifications, and reference at this meeting. The specifications list the currently installed detector area (mirror area for atmospheric Cherenkov and instrumented detector area for air shower). There is no reference at this meeting for the CACTUS telescope.

Experiment	Type	Location	Altitude	Specifications	Ref.
CACTUS	AC-Sampling	Barstow, USA	640 m	144 x 42 m ²	
CANGAROO-III	AC-Imaging	Woomera, Australia	165 m	4 x 57 m ²	[5]
HESS	AC-Imaging	Gamsberg, Namibia	1800 m	4 x 110 m ²	[6]
MAGIC	AC-Imaging	La Palma, Spain	2250 m	1 x 226 m ²	[7]
PACT	AC-Sampling	Pachmarhi, India	1075 m	25 x 4.5 m ²	[8]
SHALON	AC-Imaging	Tien Shan, Kazakhstan	3338 m	1 x 11 m ²	[9]
STACEE	AC-Sampling	Albuquerque, USA	1700 m	64 x 37 m ²	[10]
TACTIC	AC-Imaging	Mt. Abu, India	1400 m	1 x 9.5 m ²	[11]
VERITAS	AC-Imaging	Mt. Hopkins, USA	1275 m	2 x 110 m ²	[12]
Whipple	AC-Imaging	Mt. Hopkins, USA	2250 m	1 x 78 m ²	[13]
ARGO-YBJ	Air Shower	Yangbajing, Tibet	4300 m	4000 m ²	[14]
GRAPES-III	Air Shower	Ooty, India	2200 m	288 x 1 m ²	[15]
Milagro	Air Shower	Los Alamos, USA	2630 m	4800 m ²	[16]
Tibet	Air Shower	Yangbajing, Tibet	4300 m	761 x 0.5 m ²	[17]

Galactic sources: basics


- Supernova remnants = Origin of CR?
 - Energetics – OK (if 10% of E_{SN} goes to CR)
 - Maximum energy – Up to “Knee region”
 - How much of them?
 - Some evidences, which can be ascribed to HE electrons:
where are HE protons?
- Pulsar and pulsar wind nebula (plerions)
 - Crab – “The standard candle”
 - Up to a few 10GeV: pulsed+unpulsed
 - Above: unpulsed only
 - - Unpulsed: SSC (Synchrotron-Self-Compton) model
 - - Where is the cutoff?
 - - (Pulsar emission models)
 - Others?

Particle acceleration in SNR

Non-linear kinetic theory

$$t_0 = R_0/v_0; \text{ sweep up time}$$

Particle spectrum



Maximum momentum





Fig. 2. The maximum CR momentum as a function of time for the same cases as in Fig. 1.

Berezhko & Voelk, APh 1997

Berezhko & Voelk, APh 2000

Cf. Lagage and Cesarsky 1984

Nuclear gamma-ray flux from SNR




Gamma-ray emission from SNR

$n = 10 \text{ cm}^{-3}$

$n = 1 \text{ cm}^{-3}$

$n = 0.1 \text{ cm}^{-3}$




$e/p = 0.01$

0.1

1

Pulsar nebula



- A shock is formed when pulsar wind balances with ambient gas pressure, and the wind shines by synchrotron emission by thermalization




The Crab


Inner ring

=


Shock front



Optical + X-ray image



Synchrotron Self Compton




“Known” galactic sources: Supernova remnants


- **Crab** “The standard candle”
 - Well established (many observations since 1989)
- Supernova remnant **RX J1713.7-3946**
 - CANGAROO 2000/2002, H.E.S.S. 2004
- Supernova remnant **Cas A**
 - HEGRA CT system 2001
- Supernova remnant **RX J0852.0-4622**
 - CANGAROO 2005, H.E.S.S. 2005
- Supernova remnant **G0.9+0.1** [H.E.S.S. 2005]
- Supernova remnant **Vela X** [H.E.S.S. 2006, CANGAROO 2006]
- Possible SNRs: G338.3-0.0, G8.7-0.1, G18.0-0.7,
G23.3-0.3, G25.5+0.0, AX J1813-178 [H.E.S.S. 2005]
- Pulsar PSR 1706-44, Vela pulsar, SN1006
 - CANGAROO-I claims, but H.E.S.S. upper limits

SNR RX J1713.7-3946 (1)

- Detected in X-rays
- Non-thermal X-ray spectrum




Significance map



Energy spectrum

SNR RX J1713.7-3946 (2)



Hard to explain by emission from electrons (Brems, IC)

⇒ Emission from protons (π^0)?

⇒ Cosmic ray origin?

NANTEN results :

Distance ~ 1 kpc

Age ~ 1600 yr

→ $L_p \sim 10^{48} \text{ erg} \sim 0.001 L_{\text{SN}}$



(Fukui et al. PASJ 55, 2003)

SNR RX J1713.7-3946 (3)

Counter arguments

* Reimer & Pohl, A&A 390 (2002) L43

* Butt et al., Nature 418 (2002) 489



SNR RX J1713.7-3946 (4)

F. Aharonian et al.: The γ -ray supernova remnant RX J1713.7–3946

9




Fig. 9. Morphology of RX J1713.7–3946 as it appears at different energies. Shown from left to right are gamma-ray excess images with energies of $E < 0.6$ TeV, $0.6 \text{ TeV} < E < 1.4$ TeV, and $1.4 \text{ TeV} < E$. Drawn additionally as white lines are contours of significance, linearly spaced at $5, 10, 15\sigma$ (as in Fig. 7). Note the increase in the signal-to-noise ratio with increasing energy. The energy bands were chosen such that each band represents about a third of the full data set (taking events after cuts). Furthermore, all three images were smoothed with a Gaussian of $2'$, which makes them directly comparable to each other, and to Fig. 7. The resolution in each energy band is indicated in the lower left hand corner of the images; the three data subsets have comparable resolutions of $\approx 0.08^\circ$ (the resolution of the intermediate energy band is about 6% better). This might be counter-intuitive, given that at larger energies camera images get bigger and fluctuation effects become negligible thereby improving the energy and direction resolution. However, in this case that effect is compensated by the increasing mean zenith angle of the large-energy events.

SNR RX J1713.7-3946 (5)

CO

F. Aharonian et al.: The γ -ray supernova remnant RX J1713.7–3946

15




Fig. 17. Left panel: Shown are the intensity distribution of CO ($J = 1 - 0$) emission (Fukui et al. 2003) (linear colour scale in units of K km s^{-1} , truncated at a value of 23 to highlight important features), derived by integrating the CO spectra in the velocity range from -11 km s^{-1} to -3 km s^{-1} (which corresponds to 0.4 kpc to 1.5 kpc in space). Overlaid are coloured contours of the H.E.S.S. gamma-ray excess image. The levels are labelled and linearly spaced at 30, 60, and 90 counts. Note that the image is shown in Galactic coordinates. Right panel: Azimuthal profile plot, that is, number of counts as a function of the azimuthal angle, integrated in a 0.2° -wide ring covering the shell of RX J1713.7–3946 (dashed yellow circle in the left-hand panel). Plotted are the H.E.S.S. gamma-ray and the NANTEN data set.

SNR RX J1713.7-3946 (6)

Electron origin model




Fig. 19. Broadband SED of RX J1713.7–3946. The ATCA radio data and ASCA X-ray data (Hiraga 2005) for the whole SNR are indicated, along with the H.E.S.S. measurement and the EGRET upper limit. Note that the radio flux was determined in Lazendic et al. (2004) for the northwest part of the shell only and was scaled up by a factor of two here to account for the whole SNR. The synchrotron and IC spectra were modelled assuming a source distance of 1 kpc, an age T of 1000 years, a density n of 1 cm^{-3} , and a production rate of relativistic electrons by the acceleration mechanism in the form of a power law of index $\alpha = 2$ and an exponential cutoff of $E_0 = 100 \text{ TeV}$. Shown are three curves for three values of the mean magnetic field: $7 \mu\text{G}$, $9 \mu\text{G}$, and $11 \mu\text{G}$, to demonstrate the required range of the B field strength for this scenario. The electron luminosity is adopted such that the observed X-ray flux level is well matched. For the three magnetic field values the luminosity L_e is $L_e = 1.77 \times 10^{37} \text{ erg s}^{-1}$ ($7 \mu\text{G}$), $L_e = 1.14 \times 10^{37} \text{ erg s}^{-1}$ ($9 \mu\text{G}$), and $L_e = 0.81 \times 10^{37} \text{ erg s}^{-1}$ ($11 \mu\text{G}$).

Proton origin model




Fig. 20. H.E.S.S. data points plotted in an energy flux diagram. They shaded grey band is the systematic error band for this measurement (see Sect. 3.2). The black curve is the best fit of a power law with exponential cutoff to the data, extrapolated to lower energies. The dashed blue curves is the same function, but it takes the π^0 kinematics into account. The EGRET upper limit from 1 GeV to 10 GeV is plotted as red arrow.

⇒ Protons favored (?)

SNR RX J1713.7-3946 (7)




Fig. 14. The image illustrates the results of the spatially resolved spectral analysis. **Left part:** Shown in red are gamma-ray excess contours from Fig. 7, linearly spaced at 30, 60, and 90 counts. Superimposed are the 14 boxes (each $0.26^\circ \times 0.26^\circ$ in dimension) for which spectra were obtained independently. The dashed line is the 0.65° radius circle that was used to integrate events to produce a spectrum of the whole SNR. The photon index obtained from a power-law fit in each region is colour coded in bins of 0.1. The ranges of the fits to the spectra have been restricted to a maximum of 8 TeV (see Table 2). **Right part:** Plotted is the integral flux above 1 TeV against the photon index, for the 14 regions the SNR was sub-divided in. The error bars are $\pm 1\sigma$ statistical errors. Note that systematic errors of 25% on the flux and 0.1 on the photon index are to be assigned to each data point additionally.

SNR RX J0852.0-4622 [G266.2-1.2, Vela Jr.](1)





Figure 1 Rosat all-sky survey images of the Vela SNR and its surroundings. Angular resolution is 1 arcmin half-power radius; mean exposure is 993 s. The left-hand image was taken for photon energies $0.1 < E < 2.4$ keV; surface brightness increases from dark yellow to white by a factor of 500. The right-hand image is for photon energies > 1.3 keV. Most of the Vela SNR X-ray emission which dominates at low energies had disappeared. At the centre, the synchrotron nebula around the Vela pulsar remains visible as well as the SSW beam-like structure, and at the very northwest (upper right) the bright Puppis-A SNR can be seen. The new shell-type SNR RX J0852.0 – 4622 shows up in the lower left. East of RX J0852.0 – 4622 hard X-ray photons from the D/D' Vela SNR shrapnels are seen which, however, are associated with a much lower-temperature spectrum than RX J0852.0 – 4622 (ref. 14). For X-ray spectral analysis, RX J0852.0 – 4622 was divided into two

regions, one containing the bright northern limb section (l) and the other one (r) excluding the northern and southern limbs. Spectral fits were performed with either power-law models, optically thin thermal emission equilibrium models (Raymond-Smith models) or combinations of both. Solutions with a reduced $\chi^2 < 1$ for region r are obtained only with a two-temperature model with $kT_{l1} = 0.14^{+0.08}_{-0.03}$ keV, $kT_{l2} = 2.5^{+4.5}_{-0.7}$ keV. The spectrum of the northern limb can be fitted by either a simple power law with index $\alpha = -2.6^{+0.3}_{-0.4}$ or a two-temperature model with $kT_{l1} = 0.21^{+0.14}_{-0.09}$ keV, $kT_{l2} = 4.7^{+4.5}_{-0.7}$ keV. The presence of low-temperature components may partially be due to a residual, uncorrected contribution from the much softer Vela SNR. The total, absorption-corrected flux of the high-temperature components is $F_x(0.1\text{--}2.4\text{ keV}) = 3 \times 10^{-10}$ erg cm $^{-2}$ s $^{-1}$.

SNR RX J0852.0-4622 (2)



CANGAROO-II: Katagiri et al.,
ApJ, 619, (2005) L163






Fig. 2. Count map of γ -rays from the direction of RX J0852.0-4622 after background subtraction. The data are smoothed with a Gaussian ($\sigma = 0.1^\circ$) representing the angular resolution of the instrument. The point spread function (PSF) is indicated by a circle. γ -ray features smaller than the PSF should not be considered as real. The lines denote equidistant contours of smoothed ($\sigma = 0.1^\circ$) X-ray data from the ROSAT All Sky Survey, with energies restricted to above 1.3 keV. The position of the neutron star candidate AX J0851.9-4617.4 is marked with an asterisk. The axes show J2000.0 equatorial coordinates.

H.E.S.S.: Aharonian et al., AA
437, L7 (2005)

SNR RX J0852.0-4622 (3)





CANGAROO-II: Katagiri et al., ApJ, 619, (2005) L163




H.E.S.S.: Aharonian et al., AA 437, L7 (2005)

SNR RX J0852.0-4622 (4)




- Stereo (T2 & T3 wobble)
- 1,204 min. (2004 Jan/Feb)
- Off region: Vela or out of SNR
- Distance \sim 1 kpc
(NANTEN: Y.Fukui, private comm.)




SNR G0.9+0.1

- Compatible with a point source
- Position compatible with the PWN position
- Emission not consistent with the SNR shell



- Composite SNR, but no detected PSR

XMM (1.5-12 keV)
Porquet et al., 2003



H.E.S.S. Galactic plane survey



230 hr, 500 pointings

14 new TeV sources + 3
known


Scale height

$\sim 0.3\text{deg}$ RMS

\sim molecular gas



Other SNRs?




There are many SNRs!

Supernova Remnants (Green 2004)

231 SNRs

35 New SNRs by Brogan et al (2006)


Total >300 SNRs?





“New” galactic sources

- The Galactic center
 - Whipple 2004, CANGAROO 2004, H.E.S.S. 2004, MAGIC 2006
- Pulsar wind nebulae
 - **MSH15-52** [CANGAROO 2000, H.E.S.S. 2005]
 - **Vela X** [H.E.S.S. 2005, CANGAROO 2006]
- Pulsar binary **PSR 1259-63/SS2883**
 - H.E.S.S. 2005
- X-ray binary/microquasar **LS5039**
 - H.E.S.S. 2005
- UnID
 - HEGRA J2032+4130
 - H.E.S.S. J1303-631
 - New H.E.S.S. sources

The Galactic center [Sgr A*]



H.E.S.S.: Aharonian et al.,
A&A 425, L13 (2004)
/ICRC2005



CANGAROO-II (Tsuchiya
al., ApJ 606, L115, 2004)





FIG. 1.— Smoothed sky map of γ -ray candidates (background subtracted) in the direction of the Galactic Center for $\text{SIZE} \geq 300 \text{ ph. el.}$ (corresponding to an energy threshold of about 1 TeV). Overlayed are green contours (0.3 Jy beam^{-1}) of 90 cm VLA (BCD configuration) radio data (LaRosa et al. 2000). The white line shows the galactic plane.

MAGIC, ApJ 638, L101
(2006)

Dark matter signal from Sgr A*?



Neutralino-type dark matter:

Horns, Phys.Lett. B607 (2005) 225






FIG. 3. The HESS data [3] compared to the gamma-ray flux from a region of 10^{-5} sr encompassing the GC, for a $B^{(1)}$ mass of 0.8 TeV, a 5% mass splitting at the first KK level, and a boost factor b around 200 (dashed line). The solid line corresponds to a hypothetical 10 TeV WIMP with similar couplings, a total annihilation rate given by the WMAP relic density bound, and a boost factor around 1000.

Kaluza-Klein dark matter:

Bergstroem et al, Phys.Rev.Lett. 94 (2005) 131301

Pulsar wind nebula MSH15-52

PSR 1509-58



CANGAROO-I.:



Sako et al., ApJ 537, 422 (2000)

H.E.S.S.:

Aharonian et al., A&A 435, L17 (2005)

Vela X nebula

Pulsar position



θ^2 from Vela X center

- Pulsar pointing (2004 Jan/Feb)
- Stereo (T2 & T3 wobble)
- 1,311 min.
- Fisher discriminant






Fig. 1. Gaussian smoothed sky map of region surrounding Vela pulsar, showing significant emission to the south of the pulsar position, coincident with an X-ray feature seen by ROSAT (white contours). The smoothing width used is 0.09° . The contours corresponding to the strong emission close to the pulsar (position I) are truncated. The image inset in the bottom left corner indicates the size of a point source as seen by HESS, for an equivalent analysis. The solid circle represents the HESS integration region for the spectral measurement, while the dashed circle represents the field of view for the ROSAT observations. Position II is marked by a black cross.


H.E.S.S.: Aharonian et al.,

A&A 448, L43 (2006)


Pulsar binary PSR 1259-63/SS2883 (1)





3.4 year highly eccentric orbit around $\sim 10 M_{\odot}$
Be star closest approach $\sim 10^{13} \text{ cm}$ or ~ 20 stellar radii



- (i) aligned disc to the orbital plane and interaction throughout the orbit
- (ii) mis-aligned disc and interaction in the ~ 200 -day period around periastron (τ), during which the radio emission is depolarized
- (iii) mis-aligned disc and interaction in two short periods, $[(\tau - 18 \text{ d}) \sim (\tau \sim -8 \text{ d})]$ and $[(\tau + 12 \text{ d}) \sim (\tau + 22 \text{ d})]$





Pulsar binary PSR 1259-63/SS2883 (2)





X-ray binary/microquasar LS5039

- compact 4 (?) M_{\odot} object in eccentric 4 day orbit around 20-30 M_{\odot} star
- closest approach $\sim 10^{12}$ cm or ~ 2 stellar radii



What are H.E.S.S. unID sources?

- Near the Galactic plane: SNRs, pulsars?
- Hard spectra: marginally compatible with Fermi acceleration?
- Some may be identified with SNRs.
 - J1616-508, J1813-178 , J1825-137
- Old SNRs? [Yamazaki et al., astro-ph/0601704]



Aharonian et al., ApJ 636, 777 (2005)

HESS J1616-508

Young
pulsar
(69ms)






Figure 5.9: Smoothed excess map (left) of the region surrounding HESS J1616-508 (left hand source). HESS J1614-518 is also visible in this map (right hand source), along with nearby pulsars and SNRs which were considered as counterparts (smoothing radius 0.06°). The figures on the right hand side show spectrum (top) and θ^2 (bottom) plot of HESS J1616-508.


HESS J1813-178 = SNR AX J1813-178/G12.82-0.02



(a) VLA 90cm
(b) Spitzer 8 μ m /
VLA 20cm /
VLA 90cm

Less IR \leftrightarrow non-thermal


Brogan et al., ApJ 629,
L105 (2005)



INTEGRAL 20-100 keV

Ubertini et al., ApJ 629, L109 (2005)

HESS J1825-137 = PWN G18.0-0.7?



XMM: Gaensler et al., ApJ 588, 441 (2003)




Fig. 3. Spectral energy distribution of HESS J1825-137, assuming that the X-ray emission surrounding PSR B1823-13, the EGRET source 3EG J1826-1302 and the new VHE γ -ray source are related. X-ray data, indicated by lines, are taken from Gaensler et al. (2003) and are shown for the two different regions as described in the text. EGRET data (full circles) are taken from the third EGRET catalog (Hartman et al. 1999). The triangles show the HESS data from this work.

H.E.S.S.: Aharonian et al., A&A 442, L25 (2005)




Fig. 1. Excess map of the region close to PSR B1823-13 (marked with a triangle) with uncorrelated bins. The best fit centroid of the γ -ray excess is shown with error bars. The black dotted circle shows the best fit emission region size (σ_{source}) assuming a Gaussian brightness profile. The black contours denote the X-ray emission as detected by *XMM-Newton*. The 95% confidence region (dotted white line) for the position of the unidentified EGRET source 3EG J1826-1302 is also shown. The system acceptance is uniform at the 20% level in a 0.6° radius circle around HESS J1825-137.

Relation with CO/HI?







Figure 1. Multi-resolution filtered and reconstructed images of H.E.S.S. survey sources (the colour scale is arbitrary): (a) HESS J1614-518 and HESS J1616-508 (b) HESS J1640-465 (c) HESS J1813-178 (d) HESS J1825-137 (e) HESS J1837-069. Possible counterpart are also indicated. The positional confidence contours for EGRET sources are shown in dash Gray (green), ASCA sources are indicated with crosses. The SNRs are indicated as circles (red) and the molecular cloud contours are shown in white, the numbers show column intensity levels in $\text{K}.\text{kms}^{-1}$ units; the pulsars are shown as light Gray squares.

“Kookaburra” region



MOST 843MHz:
Roberts et al., ApJ
515, 712 (1999)

Fig. 2:
TeV emission as shown in Fig. 1,
together with the radio contours from
[Roberts et al. 1999](#). The TeV emission
lines up with the wings of the Kookaburra
(labeled K2/K3 and K4) and extends
north of the pulsar wind nebula
surrounding PSR J1420-6048 (K3) and
west of the Rabbit. (Preliminary)




Old SNRs: large F_{TeV}/F_X

Age $\sim 10^3$ yr

$B \sim *10 \sim *100 \mu\text{G}$

$E_{p,\text{max}} = 10^{2-3} \text{TeV}$

$E_{e,\text{max}} \sim 10 \text{TeV}$




Age $\sim 10^5$ yr

$B \sim 100 \mu\text{G}$

$E_{p,\text{max}} = 10^{1-2} \text{TeV}$

$E_{e,\text{max}} \sim 0.1 \text{TeV}$




Dim in X:
HESS unID?

Extragalactic sources: basics

- Active galactic nuclei
 - Blazars
 - Wide-band spectrum – nonthermal
 - Quasars – LBL (RBL) – HBL (XBL) sequence
 - Leptonic models
 - SSC or EC (External Compton)
 - Hadronic models
 - Proton-initiated cascades
 - Radio galaxy,...
- Gamma-ray absorption by EBL (Extragalactic Background Radiation)
 - Infrared photon field: uncertain
- Center of galaxies
 - Accumulation of dark matter??
- Extragalactic background radiation

Blazars



Beaming factor

$$\delta \equiv 1/\Gamma (1 - \beta \cos \theta) > 1$$

Observed frequency

$$\nu \propto \nu_0 \delta$$

Apparent luminosity

$$L \propto L_0 \delta^4$$

“Known” extragalactic sources

- Mrk421 ($z=0.031$)
 - First detection in 1992 [Punch et al. Nature 1992]
 - Flares in 1994, 1996, 2001, 2002-3
- Mrk501 ($z=0.034$)
 - First detection in 1995 [Quinn et al. ApJ 1996]
 - Large flares in 1997
- 1H1426+428 ($z=0.129$)
 - First detection in 2001 [Horan et al. 5th Compton 2001]
 - Flares in 2001

Multiwavelength spectra of blazars





Fig. 1. Simultaneous and non-simultaneous X-ray and TeV γ -ray energy spectra of the 4 TeV blazars with measured TeV γ -ray energy spectra. The regions show the range of values that have been observed with BeppoSAX, RXTE and Cherenkov Telescopes (from (46)).


Synchrotron self-Compton model

- Synchrotron + inverse Compton model works well
→ e^\pm origin (SSC: Synchrotron Self Compton)




Takahashi et al. ApJ 542, 2000

Synchrotron proton blazar model (1)





Muecke et al. APh 18, 2003

Synchrotron proton blazar model (2)



TeV gamma-ray absorption on EBL (1)



Mean free path for e^+e^- pair production







Figure 2: Mean free path for photon-photon pair production in the infrared-microwave background radiation. The curves correspond to those in Fig. 1 except that the effect of Lorentz Invariance violation discussed in Section 4 is shown by the long dashed curve.

TeV gamma-ray absorption on EBL (2)



EBL Spectra




AGN summary


Source	Redshift	Type	First detection	Confirmation
M87	0.004	FR I	HEGRA	H.E.S.S.
Mrk 421	0.031	BL Lac	Whipple	Many
Mrk 501	0.034	BL Lac	Whipple	Many
1ES 2344+514	0.044	BL Lac	Whipple	HEGRA
1ES 1959+650	0.047	BL Lac	7TA	Many
PKS 2005-489	0.071	BL Lac	H.E.S.S.	
PKS 2155-304	0.116	BL Lac	Durham	H.E.S.S.
H1426+428	0.129	BL Lac	Whipple	Many
H2356-309	0.165	BL Lac	H.E.S.S.	
1ES 1218+304	0.182	BL Lac	MAGIC	
1ES 1101-232	0.186	BL Lac	H.E.S.S.	

⇒ Reaching further out in redshift!

TeV gamma-ray absorption on EBL (3)



We need many samples of AGNs at various redshifts!



PG1553+113: even further away?

BL Lac, $z > 0.25$ (possibly > 0.78 ?)




Fig. 2. The energy spectrum of PG 1553+113. The horizontal error bars indicate the energy bin size. The dashed line represents the best χ^2 fit of a power law.





Fig. 3. The intrinsic photon index of PG 1553+113, with 1σ statistical error contours, versus redshift for the cases of *minimal* and *maximal* EBL absorption. An intrinsic photon index below the horizontal line ($\Gamma_{\text{int}}=1.5$) is not considered realistic. The uppermost (dashed-dotted) curve is the sum of the intrinsic photon index and 2σ statistical error for the case of minimal EBL absorption.

\Rightarrow Upper limit: $z < 0.74$

Cf. Conservative limit from non-detection of features by VLT: $z > 0.09$

TeV blazar population




Redshift distribution of blazars

C. Dermer, GLAST LAT AGN Science Group meeting, March 4, 2006


Model Fit to Blazar Redshift Distribution

Fit parameters for the FSRQs are $\Gamma = 8$ and comoving directional luminosity $l = 10^{40}$ ergs sr^{-1} s^{-1} ; EC statistics

Fit parameters for the BL Lacs are $\Gamma = 5$ and $l = 10^{42}$ ergs sr^{-1} s^{-1} ; syn/SSC statistics




Galactic diffuse gamma-rays (1)



>90% of EGRET-detected photons are diffuse!


- EGRET “GeV bump” (Hunter et al. ApJ 1997)



Galactic diffuse gamma-rays (2)

OG 2.1: Diffuse γ -ray Sources

1. Galactic Plane




Milagro [Sinnis]

~ 4.5σ detection in 3 yr data set.

Inner region:

$40\text{--}100^\circ$ in longitude, $\pm 5^\circ$ in latitude



Tibet [Ohnishi]

Flux upper limit, consistent with Milagro detection.

Diffuse gamma-rays along the plane (1)




Figure 1 | VHE γ -ray images of the Galactic Centre region. **a**, γ -ray count map; **b**, the same map after subtraction of the two dominant point sources, showing an extended band of gamma-ray emission. Axes are Galactic latitude (x) and Galactic longitude (y), units are degrees. The colour scale is in ‘events’ and is dimensionless. White contour lines indicate the density of molecular gas, traced by its CS emission. The position and size of the composite supernova remnant G 0.9+0.1 is shown with a yellow circle. The position of Sgr A* is marked with a black star. The 95% confidence region for the positions of the two unidentified EGRET sources in the region are shown as dashed green ellipses²⁰. These smoothed and acceptance-corrected images are derived from 55 hours of data consisting of dedicated observations of Sgr A*, G 0.9+0.1 and a part of the data of the HESS Galactic plane survey²¹. The excess observed along the Galactic plane consists of $\sim 3,500 \gamma$ -ray photons and has a statistical significance of 14.6 standard deviations. The absence of any residual emission at the position of the point-like γ -ray source G 0.9+0.1 demonstrates the validity of the subtraction technique. The energy threshold of the maps is 380 GeV, owing to the tight γ -ray selection cuts applied here to improve signal/noise and angular resolution. We note that the ability of HESS to map extended γ -ray emission has been demonstrated for the shell-type supernova remnants RXJ1713.7–3946 (ref. 22) and RX J0852.0–4622 (ref. 23). The white contours are evenly spaced and show velocity integrated CS line emission from ref. 11, and have been smoothed to match the angular resolution of HESS.

Diffuse gamma-rays along the plane (2)






Figure 3 | Energy distribution of Galactic cosmic rays. γ -ray flux per unit angle in the Galactic Centre region (data points), compared with the expected flux, assuming a cosmic-ray spectrum as measured in the solar neighbourhood (shaded band). The spectrum of the region $|l| < 0.8^\circ$, $|b| < 0.3^\circ$ is shown using full circles. These data can be described by a power law: $dN/dE = k[E(\text{in TeV})]^{-\Gamma}$, with $k = (1.73 \pm 0.13_{\text{stat}} \pm 0.35_{\text{sys}}) \times 10^{-8} \text{ TeV}^{-1} \text{cm}^{-2} \text{s}^{-1} \text{sr}^{-1}$ and a photon index $\Gamma = 2.29 \pm 0.07_{\text{stat}} \pm 0.02_{\text{sys}}$. The shaded box shows the range of expected π^0 -decay fluxes from this region assuming a cosmic-ray spectrum identical to that found in the solar neighbourhood and a total mass of $1.7\text{--}4.4 \times 10^7$ solar masses in the region $|l| < 0.8^\circ$, $|b| < 0.3^\circ$ estimated from CS measurements. Above 1 TeV an enhancement by a factor of 3–9 relative to this prediction is observed. Using independent mass estimates derived from submillimetre measurements²⁴, $5.3 \pm 1.0 \times 10^7$ solar masses, and from C¹⁸O measurements²⁵, $3^{+2}_{-1} \times 10^7$ solar masses, results in enhancement factors of 4–6 and 5–13, respectively (see Supplementary Information). The strongest emission away from the bright central source HESS J1745–290 occurs close to the Sgr B complex of giant molecular clouds²⁶. In a box covering this region ($0.3^\circ < l < 0.8^\circ$, $-0.3^\circ < b < 0.2^\circ$), integrated CS emission suggests a molecular target mass of $6\text{--}15 \times 10^6$ solar masses. The energy spectrum of this region is shown using open circles. The measured γ -ray flux (>1 TeV) implies a high-energy cosmic-ray density which is 4–10 times higher than the local value. Standard γ -ray selection cuts are applied here, yielding a spectral analysis threshold of 170 GeV. The spectrum of the central source HESS J1745–290 is shown for comparison (using an integration radius of 0.14°). All error bars show ± 1 standard deviation.

Diffuse source in Cygnus?


OG 2.1: Diffuse γ -ray Sources

2. Cygnus Region



Milagro [Smith]


All-sky survey – for extended sources.
Cygnus region: most luminous in N. sky.
Significance $\sim 7\sigma$.




Milagro excess (color).
EGRET diffuse model (contours).

Gamma Ray Bursts

- Ground-based experiments?
 - TeV gamma-rays (afterglow)
 - MAGIC a few per year expected
 - Air shower rate
 - Tibet-III
 - Single particle rate
 - GRAND
 - ARGO-YBJ
 - Tibet-III
 - All-sky monitor
 - Ashra
- Need fast and precise GRB alerts!



Dark matter annihilation




Signal enhancement due to ‘cusp’ structure toward the center?


“*Explosive annihilation*” by non-perturbative effect

Line

Flux ($\text{cm}^{-2}\text{sec}^{-1}$) $\Delta\Omega = 10^{-3}$



Continuum




Source counts

Source Type*	2003	2005
Pulsar Wind Nebula (e.g. Crab, MSH 15-52 ...)	1	6
Supernova Remnants (e.g. Cas-A, RXJ 1713 ...)	2	6
Binary Pulsar (B1259-63)	0	1
Micro-quasar (LS 5039)	0	1
Diffuse (Cygnus region)	0	1
AGN (e.g. Mkn 421, PKS 2155 ...)	7	11
Unidentified	2	6
TOTAL	12	32

* Includes likely associations of HESS unid sources.


→ Explosion in the number of VHE sources.

“Evolution” of the TeV gamma-ray sky




GeV vs TeV

Third EGRET catalog: Hartman et al, ApJS 123, 79 (1999)



“Evolution” in number of objects


“Kifune Plot”



Source count versus year
[T. Kifune]

Large area left unexplored at TeV


S. Funk, ICRC2005



Near future

Which Direction Should We Go?

- Low energy extension
 - Unopened window
- High sensitivity or high energy extension
 - More statistics
- Better time coverage
 - Transient sources



To lower energies...

VERITAS: VERITAS-4 by 2006



~~New site: Horseshoe canyon,
Kitt Peak, Arizona~~

Smithsonian Inst. etc.

Prototype (Aug '03)

Nov 2005: New Environmental Assessment

~~Oct 2006: Completion of Phase I:
4 telescope array~~

Then VERITAS-7 in 200X



H.E.S.S.-II

Khomas highland, Namibia (1800m a.s.l.)

THE NEXT STEP IN BOTH DIRECTIONS: HESS-II

Very Large
Cherenkov Telescope

- Lower threshold and increased energy range
 - in stand-alone mode
 - Improved sensitivity at higher energy in coincidence mode
 - Turn-over of spectra for VHE sources
 - Pulsars, Microquasars, ...
 - Unidentified sources
 - AGNs and cosmology; redshift coverage
 - GRBs
 - Dark matter

$\Phi \sim 28 \text{ m}$
 (600 m^2)
 $f \sim 35 \text{ m}$
Parabolic

$$E_{thr} \propto (\sqrt{A \epsilon / B \Omega \Delta t})^{-1}$$

A mirror area, Ω angular size
 ϵ photon detection efficiency
 B night-sky noise
 Δt integration window



To lower energies...

M. Teshima, Cherenkov2005


MAGIC-II and III

Canary island (2200m a.s.l.)



MACE

Hanle, North India (4200m a.s.l.): 2010?




To higher sensitivity...

Midsized-telescope arrays

Many conceptual designs: for example,

HE-ASTRO

Because the size of the HE-ASTRO, **~1 km²**, is much larger than the size of the Cherenkov light pool, **~10⁸ cm²**, the number of telescopes required is > 200



Coupling distance: d=80m



T. Weekes, Energy Budget in the High Energy Universe, 2006

To wider sky coverage...

Wide-field optics

GAW (Gamma Air Watch)

- Flat, single-side lens
- Lens Diameter = 2,1 m
- Focal length = 2,5 m
- $f/\# = 1.2$
- Field of view = ± 12 deg.



Mineo et al., NewHEGE3, 2005

ASHRA




M.Sasaki, Energy
Budget in the High
Energy Universe, 2006

Particle detector: HAWC

- 300m x 300m pond
- Wide FOV: $\sim 2\text{sr}$
- 100% duty cycle

- Build pond at extreme altitude (Tibet 4300m or Chile 5200m)
- Incorporate new design
 - Optical isolation between PMTs
 - Larger PMT spacing
 - Deeper PMT depth (in top layer)




~\$20M for complete detector

~60x sensitivity of Milagro – instantaneous sensitivity of Whipple over 2 sr


Crab Nebula in 30 minutes (now 1 year)

GRBs to redshift of >1 (now 0.4)



International coordination


- Monitoring of time-variable objects (e.g. blazars)
- Multiwavelength observation campaign



An example: MAGIC & H.E.S.S. campaign on Mrk 421

- Mrk 421 had the most active known period during 2004
- MAGIC has observed this object for ~14 hours in different emission states (total significance of the detection is above 40σ)
- MAGIC and HESS first combined observation of AGN Mrk421 during December 2004

Mrk421



Combined spectrum – agreement
and complementary!

Summary

- Very high energy sources may contain large varieties, including both galactic and extragalactic objects.
- Supernova remnants are confirmed to be very-high-energy particle accelerators: an important evidence of cosmic ray origin!
- TeV gamma-ray astronomy is becoming an indispensable field of astronomy.
- The “third generation” Cherenkov telescopes are working hard (and “fourth” ones are planned)— more fun!