乗鞍岳におけるミュオンの精密測定

宗像、加藤、安江、中野、石崎、宮崎(信州大理)、青木(ICRR)、小島(愛工大)

校費: 400千円 (破損パネル交換用太陽電池パネル3枚)

旅費: 150千円 (松本~乗鞍、松本~柏)

SK & Tibet空気シャワーアレイによる 10 TeV宇宙線強度の恒星時日周変動の観測

宗像、加藤、安江、中野、石崎、宮崎(信州大理)、瀧田(ICRR)

旅費(松本 $\leftrightarrow$ 柏): 50千円(SK) / 150千円(Tibet)

乗鞍ミュージオン計

2D map of 1 hour data (2006 12/14 08:30UT)

2012年度作業内容

- アンプ基板上的の高圧絶縁対策

- 太陽電池架台の再設計
(H25年度に改修予定)

- GHz帯データ通信開始
(宇宙線研)

Cycle 24 Sunspot Number Prediction (November 2012)

Hathaway/NASA/MSFC

At <<http://neutronm.bartol.udel.edu/~pyle/endplot.gif>>

Prepared
Tue, Dec 04, 2012

Log scale. Arbitrary factors are used to separate the traces.

- Doi Inthanon
- South Pole (Bares)
- South Pole
- McMurdo
- Fort Smith
- Nain
- Inuvik
- Peawanuck
- Apatity
- Thule
- Tixie Bay
- Newark
- Mawson
- Kp

10% =

9
6
3
0

チベット実験による恒星時異方性

Predicted/observed ISMF (B) orientations

Global + Midscale anisotropy model: **B** (Amenomori et al., *Astrophys. Space Sci. Trans*, **6**, 49, 2010)

Faraday rotation measure: **F** (Frisch, *Space Sci. Rev.*, **78**, 213, 1996)

ENA ribbon by IBEX: **I** (Frisch, *Space Sci. Rev.*, **78**, 213, 1996)

Starlight polarization by magnetized dust: **F1** (Astrophys. J., **760**, 106, 2012)

Miraglio "hot" regions (Abdo et al., *PRL*, **101**, 2008)

Local magnetic turbulence model

(Giacinti & Sigl, Phys. Rev. Lett., 109, 071101, 2012)

Tibet AS γ ($E_{\text{mode}} = 7$ TeV)
(Amenomori et al., Science, 314, 2006)

Ice Cube ($E_{\text{mode}} = 20$ TeV)
(Abbasi et al., ApJL, 718, 2010)

“Energy dependent smaller scale ($\leq 10^\circ$) anisotropies naturally arise from the local concrete realization of the turbulent magnetic field within the cosmic ray scattering length.”

FIG. 2 (color online). Same as Fig. 1, for boundary conditions imposed on concentric spheres around Earth with radii $R = 100, 50, 25, 10$ pc (respectively, first, second, third, and fourth columns). Upper row: $p/Z = 10^{16}$ eV. Lower row: $p/Z = 5 \times 10^{16}$ eV.

VOYAGER-1

> 0.5 MeV/nuc ions (6-Hour Avg)

Has V1 passed the heliopause?

Figure 1. Schematic diagram of global magnetic connection throughout the heliosphere. Inside the Connection Region, termination shock accelerated ACRs have access both directions along the field lines (through parallel diffusion), while GCRs do not. Beyond the disconnection boundary (dark black line), in the Disconnection Region, the opposite is true, with GCRs able to stream in along the field from the interstellar medium and no direct access from the termination shock.