


RECENT TOPICS IN TEV GAMMA-RAY ASTRONOMY

Masaki Mori


*Institute for Cosmic Ray Research,
University of Tokyo*

東京大学理学部天文教室談話会 2008年10月7日

Detection of gamma-rays



> TeV gamma-rays



$$A \sim 10^4 \text{m}^2$$




$$\Omega \sim 10^{-2} \text{sr}$$



Fermi Gamma-ray Space Telescope (2008 June-)


Atmospheric Cherenkov Telescope

Cherenkov light from gamma-ray showers
Lateral distribution & Timing distribution





Imaging Atmospheric Cherenkov Telescope

Shower profile




Focal plane image



→ Differentiation of gamma-rays from charged cosmic rays

α (image orientation angle)

Stereoscopic observation of Cherenkov light



© S.Funk, 2005


Angular resolution

$0.25\text{deg} \rightarrow 0.1\text{ deg}$

Energy resolution

$30\% \rightarrow 15\%$

Better S/N (no local muons)



Comparison of detection methods

Base	Satellite	Ground	Ground
Gamma-ray detection	Direct (pair creation)	Indirect (atmospheric Cherenkov)	Indirect (shower array)
Energy	< 30 GeV (→ 100 GeV)	>100 GeV (→ 50 GeV)	>3 TeV (→ 1 TeV)
Pros	High S/N Large FOV	Large area Good $\Delta\theta$	24hr operation Large FOV
Cons	Small area High cost	Low S/N (CR bkgd.) <i>(but imaging overcomes this!)</i> Small FOV	Low S/N (CR bkgd.) Moderate $\Delta\theta$

VHE Experimental World

MILAGRO



STACEE



MAGIC



TIBET



MILAGRO

STACEE

VERITAS

MAGIC

TACTIC

TIBET

ARGO-YBJ

PACT

GRAPES

2007

VERITAS



HESS

HESS



TACTIC



CANGAROO III

CANGAROO




Crab: the first TeV source


ガンマ
線信号



Whipple 10m telescope




37ch imaging camera



82hrs, $0.24\gamma/\text{min}$

Crab: 2008


Clock



Crab Signal as seen in VERITAS in real time
during commissioning


TeV skymap

1999




TeV skymap

2001




TeV skymap

2003




TeV skymap

2005



TeV skymap


2007




Increase of TeV sources

“Kifune plot”

Class	2003	2005	2007
PWN (Pulsar Wind Nebulae)	1	6	18
SNR (Supernova remnants)	2	3	7
Binary	0	2	4
Diffuse	0	2	2
AGN (Active Galactic Nuclei)	7	11	19
UnId (Unidentified sources)	2	6	21
Total	12	33	71!





TeV sources classified



* Borders for SNR/PWN/UnID are vague...


H.E.S.S. Galactic survey




Survey region was extended in the years 2005 - 2007

Hoppe 269


$-85^\circ < l < 60^\circ$ $-2.5^\circ < b < 2.5^\circ$




H.E.S.S. Galactic survey



Supernova remnants


- 
- ▣ Long considered to be primary source for Galactic cosmic rays
 - ▣ Pros:
 - Energetic enough (10% of SN explosion energy)
 - Size of object is large enough ($R \gg r_g$)
 - Many SNRs are bright radio sources: at least electrons are accelerated!
 - ▣ Cons:
 - Magnetic fields too low to go beyond 10^{14}eV
 - Additional problem: adiabatic losses

Shell SNRs seen at TeV



Comparable
to PSF (0.09°)


J. Albert et al., A&A 474,
934 (2007)




Good keV-TeV
Correlation!

T.Tanaka, PhD thesis, 2007

Accelerated particles and gamma-ray spectrum



©A.Bamba




SNR spectrum


Hard power-law + cutoff (?): $\sim E^{-2} \exp(-E/E_{\max})$

Electron model


RX J1713.7-3946



Proton model



RX J0842.0-4622



NO definitive answer
for accelerated particles!

Identification of particles is not easy






FIG. 12.—Pion-decay and IC emission for a range of n_H and B_0 . In the top panel, the heavy curves are pion decay, the light curves are IC, and $\epsilon_{\text{rel}} = 36\%$ and $B_0 = 15 \mu\text{G}$ in all cases. The strong dependence of pion decay on ambient density n_H is evident. The middle panel shows IC, and the bottom panel shows pion decay for $n_H = 0.1 \text{ cm}^{-3}$, with B_0 varying from $3 \mu\text{G}$ (solid curves) to $15 \mu\text{G}$ (dashed curves) to $60 \mu\text{G}$ (dotted curves). For comparison to the π^0 , we show in the bottom panel the IC emission for $B_0 = 60 \mu\text{G}$ (light dotted curve). The particle distributions producing the emission in the bottom two panels are those shown in the top panel of Fig. 11.

Difficult in the GeV-TeV region if magnetic field is strong!

Magnetic field in SNR

RX J1713.7-3946 by Chandra




Variation in ~ 1 yr time scale
 $\rightarrow$ Need > 1 mG ! (locally)
 $\rightarrow$ Protons produce TeV gamma-rays!?

Counter arguments: Y. Butt et al. , arXiv:0801.4954

SNR	Dist kpc	V_s km s^{-1}	n_0 cm^{-3}	width "	B_{loss} μG	B_{diff} μG
Cas A	3.4	5200	3	0.5	249	299
Kepler	4.8	5300	0.35	1.5	97	113
Tycho	2.4	4500	0.3	2	113	165
SN1006	2.2	4300	0.1	20	30	39
RCW86	2.5	3500	0.1	45	24	14


SNRs interacting with clouds?

IC443






[MAGIC, Albert et al. , ApJ 664, L87,2007]

W28




[H.E.S.S., Aharonian et al. , A&A, in press]



Evidence of proton acceleration?

SN1006

Chandra and H.E.S.S. Morphology




➔ Chandra map smoothed by the H.E.S.S. PSF and oversampled with 0.1°


➔ H.E.S.S. map with Chandra contours



- Independent analysis methods confirm the detection of the NE rim of SN 1006 at about 6 sigma in the region pre-defined in the 2004 H.E.S.S. upper limit paper
- Compatibility with non-thermal X-ray morphology very good
- Given the flux level of $\sim 1\%$ Crab, both leptonic and hadronic scenarios are reasonable




Pulsar nebulae




- Major group in Galactic TeV sources
 - 18/71 by Hinton (2007ICRC)
 - Associated with relatively young ($<10^5$ years) and large spin-down pulsars
- Extended $O(10\text{pc})$, displaced from pulsars
- Gamma-rays via inverse Compton by electrons?


Pulsar nebula: energy-dependent morphology



- **HESS J1825-137 associated with energetic pulsar**
- **Spectral steepening seen away from the pulsar**
- **Very likely this is evidence for cooling of electrons in the Nebula**
 - Seen in several *X-ray* PWN
- **A first in gamma-ray astronomy!**




Simulations of Pulsar Wind Nebula




Displacement due to pulsar kick ?


More pulsar nebulae...




Djannati-Atai et al., ICRC2007




Hoppe et al., ICRC2007



Carrigan et al., ICRC2007



Komin et al., ICRC2007



Aharonian et al. ApJ 777 (2007)

Pulsar nebulae and spin-down luminosity




Figure 3: **Top:** $P - \dot{P}$ diagram for pulsars: all ATNF pulsars (black), with detected X-ray PWN (brown), with a known corresponding SNR (blue), potentially associated to an EGRET source (green), associated to a H.E.S.S. VHE PWN (red). **Bottom:** Energy output for the selections used at the top.

Pulsed emission from Crab






Figure 4: Crab pulsar spectral cutoff. The black flux point indicates the energy range and the statistical error of our flux measurement, assuming an exponential and a super-exponential cut-off. The black crosses on the left represent flux measurements from EGRET (15). The arrows on the right denote upper limits from various previous experiments. The EGRET spectrum has been extrapolated with an exponential (red) and a super-exponential (blue) shape. The coloured bands illustrate the combined statistical and systematic errors and denote the total uncertainty of our measurement, by which the flux point can move to higher or lower energies.


→ Favors *outer-gap* model rather than *polar-cap* model



Gamma-ray binaries

LSI +61 303 (VERITAS/MAGIC)


$P_{\text{orb}} \sim 26.5 \text{ day}$



J. Albert et al., Science 312, 1771 (2006)
V.A. Acciari et al., arXiv:0802.2363

LS 5039 (H.E.S.S.)



$P_{\text{orb}} \sim 3.9 \text{ day}$




F. Aharonian et al., A&A 460 (2006) 743

Gamma-ray binary : Cyg X-1

- ▣ Black hole binary: $M_{\text{BH}} \sim 21 M_{\odot}$,
 $M_{\star} \sim 30 M_{\odot}$
- ▣ Relativistic jet $v > 0.6c$: “microquasar”
- ▣ MAGIC 40hr obs.
- ▣ 4.9σ seen in one 79 min. time slice
- ▣ Estimated significance: 4.1σ after correction for statistical trials



Emission from binaries





Microquasar: particles (electrons or hadrons) are accelerated in a jet
Bosch-Ramon et al. (2006), Romero et al. (2007)

γ -rays produced in the shock where the wind of the young pulsar and the wind of the Be star collide
Dubus (2006), Dhawan et al. (2006)

Stellar cluster Westerlund 2



- Young open stellar cluster
 - Dozen O-stars
 - Two Wolf-Rayet stars ($\sim 80M_{\odot}$ each)!
- Extended gamma-ray emission covering (but offset from) Westerlund 2 by HESS
- Due to collective effects of stellar winds in the cluster?
- A new source class?



Galactic center $\approx$ Sgr A*

HESS data 2003-2004 towards galactic centre. (We await 2005-6 data eagerly...)

MAGIC (2006) data agree with HESS




Energy spectrum is *not* consistent with dark matter annihilation signal!

Steady (time-independent) spectrum, pointlike within HESS angular resolution, could be Moore cusp instead of NFW?



But: Probably too high energy (and wrong shape of spectrum) for WIMP annihilation explanation

Shape of these curves uncertain, depends on QED corrections and fragmentation of 5-10 TeV jets. LHC should give important input here.


Galactic center ridge



Spectrum is harder than CR spectrum!



Unidentified HESS sources




F. Aharonian et al., A&A 442, 1 (2005)

Two types:


- 1) No compelling counterparts
- 2) Dark in other wavelengths

TeV-GeV relation?

Coincident sources




EGRET source	VHE γ -ray source	Potential Counterpart
Within the H.E.S.S. GPS		
3EG J1639-4702	HESS J1640-465	G338.3-0.0 (SNR/PWN)
3EG J1744-3011	HESS J1745-303	
3EG J1800-2338	HESS J1801-233	W28 (SNR)
3EG J1826-1302	HESS J1825-137	G18.0-0.7 (PWN)
3EG J1824-1514	HESS J1826-148	LS 5039 (Binary)
Outside the H.E.S.S. GPS		
3EG J0241+6103	MAGIC J0240+613	LSI +61 303 (Binary)
3EG J0617+2238	MAGIC J0616+225	IC443 (SNR/PWN)
3EG J0634+0521	HESS J0632+058	Monoceros
3EG J1420-6038	HESS J1420-607	Kookaburra (PWN)




Extended H.E.S.S. survey

Acceptance-corrected Exposure




Extended H.E.S.S. GPS

- $-85^\circ < l < 60^\circ$
- $-3^\circ < b < 3^\circ$
- Scan mode: 400 h
- Detected **50+ Galactic sources** of VHE gamma-rays
- ICRC 2007, DPG 2008, Gamma08



H.E.S.S. galactic plane survey

H.E.S.S. GPS Significance Map




- HESS J1714-385 (SNR CTB 37A)
- HESS J1713-381 (SNR CTB 37B)
- HESS J1442-623 (SNR RCW 86)
- HESS J1356-654 (Offset PWN)
- HESS J1503-582 (FVW?)
- HESS J1848-018 (SFR W 43, WR 121a?)
- HESS J1849-000 (PWN)

Extragalactic TeV sources

Name	Discovered	Year	z	Contributions
M 87	HEGRA	2003	0.004	VERITAS-Colin, HESS-Beilicke, MAGIC-
Mrk 421	Whipple	1992	0.031	MILAGRO-Smith, VERITAS-Fegan, +
Mrk 501	Whipple	1996	0.034	TACTIC-Godambe, MAGIC-Paneque, +
1ES 2344+514	Whipple	1998	0.044	MAGIC-Wagner
→ Mrk 180	MAGIC	2006	0.046	MAGIC-Mazin
1ES 1959+650	TA	2002	0.047	MAGIC-Hayashida
→ BL Lac	MAGIC	2006	0.069	MAGIC-Hayashida
→ PKS 0548-322	HESS	2006	0.069	HESS-Superina
PKS 2005-489	HESS	2005	0.071	HESS-Costamante
PKS 2155-304	Durham	1999	0.116	HESS-Punch, CANGAROO-Sakamoto, +
H 1426+428	Whipple	2002	0.129	VERITAS-Krawczynski
→ 1ES 0229+200	HESS	2007	0.140	HESS-Raue
H 2356-309	HESS	2005	0.165	HESS-Costamante
1ES 1218+304	MAGIC	2005	0.182	MAGIC-Hayashida
1ES 1101-232	HESS	2005	0.186	HESS-Puelhofer
→ 1ES 0347-121	HESS	2007	0.188	HESS-Raue
→ 1ES 1011+496	MAGIC	2007	0.212	MAGIC-Mazin
→ PG 1553+113	HESS/MAGIC	2005	?	MAGIC-Wagner, HESS-Benbow
→ 3C 279	MAGIC	2007	0.536	MAGIC-Teshima


8 new AGN

Emission from AGNs



Electron model



Fossati et al. 1998



Proton model

Muecke et al. APh 18, 2003

Fast time variation



M. Hayashida, ICRC 2007


Fast variation $\leftrightarrow$ Acceleration site & mechanism

Absorption by IR background


Observed spectrum is affected by
integralactic absorption!

$$\gamma_{\text{TeV}} + \gamma_{\text{IR}} \rightarrow e^+ + e^-$$

Mean free path for e^+e^- pair production




Protheroe & Meyer, Phys.Lett. B493 (2000) 1



Jim Hinton, rapporteur talk, ICRC 2007

We cannot discriminate
source spectrum and
intergalactic absorption!


Unfolding source spectra



Assume not
harder than $E^{-1.5}$

Some models
can be rejected

Background IR intensity limited by TeV observations



Upper limits: fluctuation/direct measurements
 Lower limits: source counts

3C279 at $z=0.538$




Fig. 1. Light curves. MAGIC (top) and optical R-band data (bottom) obtained for 3C 279 from February to March 2006. The long-term baseline for the optical flux is at 3 mJy.





Fig. 2. Spectrum of 3C 279 measured by MAGIC. The gray area includes the combined statistical (1σ) and systematic errors, and underlines the marginal significance of detections at high energy. The dotted line shows compatibility of the measured spectrum with a power law of photon index $\alpha = 4.1$. The blue and red triangles are measurements corrected on the basis of the two models for EBL density, discussed in the text.




Redshift distribution of blazars


Model Fit to Blazar Redshift Distribution

Fit parameters for the FSRQs are $\Gamma = 8$ and comoving directional luminosity $l = 10^{40} \text{ ergs sr}^{-1} \text{ s}^{-1}$; EC statistics

Fit parameters for the BL Lacs are $\Gamma = 5$ and $l = 10^{42} \text{ ergs sr}^{-1} \text{ s}^{-1}$; syn/SSC statistics



Redshift distribution of blazars









10xTeV AGN


More AGNs as energy threshold goes lower!

Clusters of galaxies: upper limits

H.E.S.S.: Domainko et al., ICRC2007




Gamma-ray skymap




(background: Fermi Gamma-ray Space Telescope Firstlight data [>100 MeV])

Under construction

H.E.S.S. II




MAGIC II





Future projects


CTA (Cherenkov Telescope Array): EU++



AGIS (Advanced Gamma-ray Imaging System): USA++



log N-log S relation



Data: H.E.S.S. catalog

× 2 (North/South)
↓
1000 TeV sources
if mCrab!

Summary


- ▣ High energy window of the Universe is now open!
 - Additional 2-3 decades of the photon spectrum
 - Wider variety of sources than expected
 - *Cosmic accelerators are ubiquitous!*
 - Much work left to understand their physics
 - Also: cosmology, fundamental physics
- ▣ Hoping to detect other class of sources...
 - Pulsars
 - Star-forming galaxies, mergers
 - Dwarf galaxies and dark matter
 - Ultraluminous IR galaxies
 - Clusters of galaxies
 - Gamma-ray bursts

Thank you!



CANGAROO-III telescopes, Woomera, South Australia

Gamma-ray signal from DM annihilation at the Galactic center



gamhalo_egret_big

Indirect detection through γ -rays.
Three types of signal:

- Continuous from π^0 , K^0 , ... decays and
- Monoenergetic line and
- Internal bremsstrahlung from QED process.

Enhanced flux possible thanks to halo density profile and substructure (as predicted by CDM)

Good spectral signatures!

Unfortunately, large uncertainties in the predictions of absolute rates

Gamma-rays

