

## **Evidence for Two Energetic Particle Sources at the Termination Shock**

E. C. Stone<sup>a</sup>, A. C. Cummings<sup>a</sup>, F. B. McDonald<sup>b</sup>, B. C. Heikkila<sup>c</sup>, N. Lal<sup>c</sup> and W. R. Webber<sup>d</sup>

(a) *California Institute of Technology, Pasadena, CA 91125, USA*

(b) *University of Maryland, College Park, MD 20742, USA*

(c) *NASA/Goddard Space Flight Center, Greenbelt, MD 20771, USA*

(d) *New Mexico State University, Las Cruces, NM 88003, USA*

Presenter: E. C. Stone (ecs@srl.caltech.edu), usa-stone-E-abs3-sh31-poster

During the period of unusual enhancements in the intensities of low energy ions observed by Voyager 1 beyond 85 AU, the energy spectra appear to have two components. Upstream of the termination shock, low energy ions (<10 MeV/nucleon) stream along the magnetic field and have significant variations in intensity and spectral slope. Their power law spectrum often shows no evidence of modulation at the lowest energies. This suggests a nearby source region of termination shock particles (TSPs) connected to Voyager along the interplanetary magnetic field. At intermediate energies (~10 to 60 MeV/nucleon), the helium spectrum is relatively steady and consistent with that of anomalous cosmic rays (ACRs) with modulation increasing with decreasing energy. This indicates that the ACR source is a remote region of the termination shock not connected to Voyager along the magnetic field. This distinction between the TSP and ACR sources was confirmed when Voyager 1 crossed the termination shock at 94 AU.

### **1. Acknowledgements**

This work was supported by NASA under contract NAS7-03001.

