

Comparison of UHE Composition Measurements by Fly's Eye, HiRes-prototype/MIA and Stereo HiRes Experiments

P. Sokolsky and the HiRes Collaboration

High Energy Astrophysics Institute, University of Utah, Salt Lake City, Utah 84112, USA

Presenter: C.C.H. Jui (ps@cosmic.utah.edu), usa-sokolsky-P-abs1-he14-oral

We compare the elongation rate and Xmax distributions for three air-fluorescence experiments: Stereo Fly's Eye, HiRes/MIA, and HiRes. A shift of 13 gm/cm² of the stereo Fly's Eye data, well within the quoted systematic errors, brings the elongation rates and the Xmax distributions of all three experiments into reasonable agreement. We explore the implications of this combined data set ranging from 10¹⁷ eV to near 10²⁰ eV.

1. Introduction

There have now been three fluorescence-based experiments which have reported results on the cosmic ray composition using the shower maximum (Xmax) method. These are the Stereo Fly's Eye [1], the HiRes Prototype/MIA[2] and the HiRes[3] experiments. The Fly's Eye collaboration reported on data taken with the Stereo Fly's Eye configuration. In this experiment, two fluorescence detectors were positioned 3.5 km apart. The effective energy range extended from 10¹⁷ eV to 3×10¹⁹ eV. This experiment utilized optics that resulted in a 5×5 degree pixel size, and the geometry of the event was determined by the intersection of the two event-detector planes, with resulting resolution in Xmax of about 45 gm/cm². Both Fly's Eye detectors had mirrors that covered an elevation angle range from 0 to 90 degrees, though FE II had only partial azimuth coverage.

The HiRes Prototype/MIA experiment was a hybrid fluorescence - muon array detector. The fluorescence detector, a fourteen mirror prototype of the current HiRes detector had a 1×1 degree pixel size and was located 3.5 km from the center of the MIA muon array. The muon array sampled a part of muon lateral distribution. The effective area of this "array" was thus set by the muon lateral distribution function and was roughly a 3 km radius centered on MIA. The smaller pixel size of the HiRes detector allowed triggering to lower energies (approaching 5×10¹⁶ eV) while the effective area of MIA limited the energy reach to just above 10¹⁸ eV. The Xmax resolution was 45 gm/cm². The geometry of the event was determined by using the HiRes event-detector plane and timing information from the fluorescence detector and the MIA ground array.

The HiRes stereo detector consists of two fluorescence detectors separated by 12 km. With 1×1 deg pixel size, the Xmax resolution is 30 gm/cm². The large separation between detectors and the limited elevation angle range (3 to 32 degrees) limits the energy range to > 10¹⁸ eV with the statistical reach approaching 10²⁰ eV.

2. Previous Results

All three experiments have published results on the cosmic ray composition in various energy ranges. Figure 1 shows the elongation rate results, together with expectations from various hadronic models for a purely

protonic and a purely Fe composition. In each case, there is evidence for a transition from a predominantly heavy composition, near 10^{17} eV to a much lighter composition at higher energies. While the qualitative picture is consistent between the experiments, the details appear to be different, with the Stereo Fly's Eye results indicating a more gradual transition while the combined HiRes Prototype/MIA and HiRes results indicate a transition to a light composition occurring more abruptly near 10^{18} eV. The fits to the elongation rates are also somewhat different, with the Stereo Fly's Eye quoting an overall elongation rate of 78.9 ± 3.0 gm/cm²/decade, while the HiRes Prototype/MIA experiment quotes 93.0 ± 8.5 gm/cm²/decade below 10^{18} eV and 54.5 ± 6.5 gm/cm²/dec above this value.

The elongation rate figures only show the error on the mean of the Xmax distribution in each energy bin. All three experiments quote systematic errors between 20 to 25 gm/cm² on the absolute position of Xmax. These are largely dominated by mirror Čerenkov light subtraction. Differences in elongation rate slopes between different experiments can in principle also be produced by non-linear energy response. The Fly's Eye Stereo experiment shows a significant ankle structure with a minimum near 3×10^{18} eV. The HiRes detector sees a very similar ankle structure with a minimum at the same energy. The flux normalization between the two spectra is also similar, reinforcing the evidence that the energy scales between the two experiments are very similar, at least in this energy range.




Figure 1. Elongation rate for Stereo Fly's Eye (left) and HiRes/MIA and HiRes (right) experiments

3. Applying a Systematic Shift

The resolution in Xmax tends to improve with increasing energy as more of the extensive shower is measured. Čerenkov subtraction effects tend to be minimized for the higher energy events because they are usually viewed at large angles relative to the shower axis. It is thus reasonable to assume that the underlying systematic shifts will be most cleanly detected at the higher energies. In this paper, we take the HiRes $>10^{18}$ eV Xmax distribution and the Stereo Fly's Eye $>10^{18}$ eV Xmax distribution and assume that the shift in the mean Xmax, averaged over the energies is representative of the overall systematic shift between these two experiments. We then apply the same systematic shift to the lower energy Stereo Fly's Eye data and check for consistency with the

HiRes Prototype/MIA data. The mean X_{\max} shift is 13 gm/cm^2 , well within the quoted systematic errors on both experiments. Fig 2 shows the elongation rate data for all three experiments with all the Fly's Eye stereo points shifted upwards by 13 gm/cm^2 . The data over the entire energy range are quite consistent, even within the statistical errors on the mean. While the HiRes elongation rate above 10^{18} eV is consistent with the shifted Stereo Fly's Eye points above this energy, the data can also be fit over the entire energy range to a single elongation rate slope of $68.5 \pm 1.5 \text{ gm/cm}^2$. Combining all the data after a simple shift consistent with systematic errors shows both the good consistency in the data itself and the importance of a wide dynamic range in energy in interpreting apparent changes in the elongation rate.

4. Comparison of X_{\max} Distributions

The fluctuations of showers in the atmosphere are also an important indication of the cosmic ray composition. From very simple superposition arguments, one expects EAS produced by heavy nuclei such as Fe to have smaller fluctuations and occur higher in the atmosphere than EAS produced by protons. Detailed comparisons to predictions have been made for each of the experiments. Here we examine the consistency of the data for the three experiments.

We ask if the Stereo Fly's Eye stereo, HiRes/MIA and HiRes experiments are consistent in the reported widths of the measured X_{\max} distributions. For historical reasons, we consider the X_{\max} distributions in energy bins of $<3 \times 10^{17} \text{ eV}$ (HiRes prototype/MIA), $3 \text{ to } 5 \times 10^{17} \text{ eV}$ (Stereo Fly's Eye and HiRes prototype/MIA), $5 \times 10^{17} \text{ to } 1 \times 10^{18} \text{ eV}$ (Fly's Eye Stereo) and $>1 \times 10^{18} \text{ eV}$ (Stereo Fly's Eye Stereo and HiRes). Fig 2 and Fig 3 show the normalized X_{\max} distributions for the two energy bins where we have data from two different experiments.

Note that we have shifted the X_{\max} for all Stereo Fly's Eye events by the canonical 13 gm/cm^2 , as above. The X_{\max} distributions are quite consistent. Note that the quoted X_{\max} resolutions for Stereo Fly's eye and HiRes/MIA are quite similar, with similar resolutions. HiRes has significantly better resolution than Stereo Fly's Eye above 10^{18} eV , but the fluctuations are likely dominated by the large intrinsic proton fluctuations and hence the difference in detector resolution is not noticeable. Fig 3 shows the normalized X_{\max} distributions for all the energy bins in one figure (for clarity we have only used the Stereo Fly's Eye and the HiRes distribution in the $3\text{-}5 \times 10^{17} \text{ eV}$ and $>10^{18} \text{ eV}$ energy bins respectively.) It is clear that the X_{\max} distributions broaden with increasing energy, as expected for a transition from a heavy to a light composition.

5. Conclusion

While additional small systematic shifts from, for example, Čerenkov subtraction, may be present at energies lower than 10^{18} eV , a simple 13 gm/cm^2 shift of the Stereo Fly's Eye data brings all X_{\max} distributions and the elongation rate into reasonable agreement. The break in the elongation rate, indicating a transition from a region of changing composition to a light composition appears to lie somewhere between 10^{18} and $3 \times 10^{18} \text{ eV}$ in the combined data. It will be important to elucidate this with higher precision in a single experiment with a broad dynamic range in energy, since the interpretation of the ankle structure as either a dip due to e^+e^- pair production of extragalactic protons off the 2.7 degree BB radiation or a transition from a galactic to an extragalactic spectrum very much depends on it[4].

References

- [1] D.J. Bird et. al., Ap. J. 424, 491 (1994); D. J. Bird et. al., Phys. Rev. D 47, 1919 (1993).
- [2] T. Abu-Zayyad et. al., Phys. Rev. Lett. 84, 4276 (2000); T. Abu-Zayyad et al., Ap. J. 557, 686(2001).
- [3] T. Abu-Zayyad et al., Ap. J. 622, 910 (2005).
- [4] V. Berezhinsky, A. Z. Gazizov and S. I. Grigorieva, Phys.Lett. B612 (2005) 147-153




Figure 2 . Left: Elongation rate. Open circles – Stereo Fly’s Eye; Filled circles – HiRes ; Triangles – HiRes/MIA. Stereo Fly’s Eye data has been shifted by 13 gm/cm². Errors are errors on the mean. Right: Xmax Distribution in 3×10¹⁷ to 5×10¹⁷ eV bin. Triangles: Stereo Fly’s Eye Stereo; Circles: HiRes/MIA.




Figure 3. Right: Xmax distribution in >10¹⁸ eV energy bin. Triangles: Stereo Fly’s Eye; Circles: HiRes. Left: Sequence of Xmax distributions in three increasing energy bins 3-5×10¹⁷, 5-10×10¹⁷ and > 10¹⁸eV.