

GW 150914-like Black Holes as Galactic High Energy Sources

Kunihito Ioka

(Center for Gravitational Physics,
YITP, Kyoto U.)

KI, Matsumoto, Teraki,
Kashiya & Murase in prep.



We Did It!

@YITP
midnight




Champagne



GW150914

**1st direct
detection
BH-BH**
 $36M_{\odot} + 29M_{\odot}$
 $L \sim 200M_{\odot}c^2/s$
 $\sim 10^{-3} c^5/G$




30-350Hz bandpass
First at L1
 $6.9 + 0.5 - 0.4$ ms
later at H1

The Most Luminous in the Universe

- Total radiated mass
 $M \sim 3.0+0.5-0.5 M_{\odot}$
- Energy $E=Mc^2 \sim 6 \times 10^{54}$ erg
- $L_{\text{peak}} \sim 3.6+0.5-0.4 \times 10^{56}$ erg/s
 $\sim 200+30-20 M_{\odot}c^2/\text{s}$
 $\sim 10^{-3} c^5/G$
- $> L_{\text{Gamma-Ray Bursts}} \sim 10^{50-54}$ erg/s

Solar mass is radiated within ~ 0.01 sec!

GW151226



2nd event

BH-BH

$14.2M_{\odot}$

$+7.5M_{\odot}$

$L \sim 170M_{\odot}c^2/s$

$a_{1 \text{ or } 2} > 0.2$

LVT151012

$R_{\text{GW}} \sim 9-240$


events

$\text{Gpc}^{-3} \text{ yr}^{-1}$

Galactic BHs

$70 \text{ Gpc}^{-3} \text{ yr}^{-1} \div 0.01 \text{ galaxy Mpc}^{-3} \times 10^{10} \text{ yr}$
 $\sim 70000 \text{ Merged BHs/galaxy}$

Galactic BHs



$70 \text{ Gpc}^{-3} \text{ yr}^{-1} \div 0.01 \text{ galaxy Mpc}^{-3} \times 10^{10} \text{ yr}$
 $\sim 70000 \text{ Merged BHs/galaxy}$

Old Problem

- Eddington 20's
- Hoyle & Lyttleton 39
- Bondi & Hoyle 44
- Bondi 52
- Zel'dovich 64
- Salpeter 64
- Lynden-Bell 69
- Shvartsman 71
- Michel 72
- Shapiro 73
- Shakura & Sunyaev 73
- Meszaros 75
- Ipser & Price 77, 82, 83
- Grindlay+ 78
- Carr 79
- McDowell 85
- Campana & Pardi 93
- Heckler & Kolb 96
- Fujita+ 98
- Popov & Prokhorov 98
- Armitage & Natarajan 99
- Agol & Kamionkowski 02
- Chisholm+ 03
- Barkov+ 12
- Motch & Pakull 12
- Fender+ 13

GWs put a lower limit on #(spinning BHs)

Spin Energy

$$E_{\text{spin}} = \left(1 - \sqrt{\frac{1 + \sqrt{1 - a_*^2}}{2}} \right) Mc^2$$

$$\cong 7\% \times Mc^2 \sim 1 \times 10^{54} \text{ erg} \left(\frac{M}{10M_{\odot}} \right)$$


$$E_{\text{tot}} \sim N_{\text{BH}} E_{\text{spin}} \sim 7 \times 10^4 \text{ BHs} \times 1 \times 10^{54} \text{ erg}$$

$$\sim 9 \times 10^{58} \text{ erg}$$

$$\sim \frac{10^{10} \text{ yr}}{100 \text{ yr}} \text{ supernovae}$$

**Comparable to
supernovae
ever happened!**

Blandford-Znajek Effect



Blandford & Znajek 77
Komissarov & Barkov 07
Tchekhovskoy+ 11
Armitage & Natarajan 99
Barkov, Khangulyan & Popov 12

Bondi Accretion

$$r_B \sim \frac{GM}{V^2} \sim 1 \times 10^{15} \text{ cm}$$


$$\times \left(\frac{M}{10 M_\odot} \right) \left(\frac{V}{10 \text{ km s}^{-1}} \right)^{-2}$$

$$V = \sqrt{c_s^2 + v^2 + v_{\text{GW}}^2}$$


$$\dot{M} \sim 4\pi r_B^2 V \rho$$

$$\sim 5 \times 10^{35} \text{ erg s}^{-1} \left(\frac{n}{10 \text{ cm}^{-3}} \right) \left(\frac{M}{10 M_\odot} \right)^2 \left(\frac{V}{10 \text{ km s}^{-1}} \right)^{-3}$$

$$\sim 4 \times 10^{-4} \dot{M}_{\text{Edd}} \left(\frac{n}{10 \text{ cm}^{-3}} \right) \left(\frac{M}{10 M_\odot} \right) \left(\frac{V}{10 \text{ km s}^{-1}} \right)^{-3}$$



Disk



$$\frac{\delta\rho}{\rho} \sim \left(\frac{L}{10^{20} \text{ cm}} \right)^{1/3}$$

$$\ell \sim \frac{1}{4} \frac{\delta\rho}{\rho} v r_B$$

$$\ell_K \sim \sqrt{GM r_{\text{disk}}}$$

$$\Rightarrow \frac{r_{\text{disk}}}{r_S} \sim 2 \times 10^4 \left(\frac{M}{10 M_8} \right)^{2/3} \left(\frac{V}{10 \text{ km s}^{-1}} \right)^{-10/3}$$

$\Rightarrow$ Disk formation

Disk axis fluctuates with $\Delta t \sim \frac{2r_B}{V} \sim 80 \text{ yr}$

Fujita+ 98

Agol & Kamionkowski 02

ADAF (Hot, Thick Disk)

Advection Dominated Accretion Flow

$$L_{\text{disk}} \sim \alpha_{\text{QED}} \frac{m_e}{m_p} \frac{\dot{M} c^2}{L_{\text{Edd}}} \dot{M} c^2 \sim 10^{-13} L_{\text{Edd}} \left(\frac{\dot{M} c^2 / L_{\text{Edd}}}{10^{-4}} \right)^2 \sim 10^{27} \text{ erg s}^{-1}$$

$$\frac{10^{27} \text{ erg s}^{-1}}{4\pi (\text{kpc})^2} \sim 10^{-17} \text{ erg s}^{-1} \text{ cm}^{-2}$$

Nearby BH disks might be observable


But Hirotani-san's talk

Ichimaru 77, Narayan & Yi 94

Fujita, Inoue, Nakamura, Manmoto, Nakamura 98

Matsumoto et al. in preparation

Magnetic Field



$$B \propto R^{-2}$$

$$3\mu\text{G} \rightarrow 3 \times 10^{10} \text{ G}$$

But B is saturated
so that $p_B < p_{\text{disk}}$

$$p_B = \frac{B^2}{8\pi}$$

$$p_a = \frac{GM\Sigma}{r^2} \sim \frac{GM\dot{M}}{2\pi r^3 v_r}$$

**Magnetically
Arrested
Disk (MAD)**


Bisnovatyi-Kogan
& Ruzmaikin 76
Narayan+ 03

$$B_H \sim \sqrt{\frac{4GM\dot{M}}{r^3 v_r}} \bigg|_{r=r_H} \sim 4 \times 10^7 \text{ G} \left(\frac{n}{10 \text{ cm}^{-3}} \right)^{1/2} \left(\frac{V}{10 \text{ km s}^{-1}} \right)^{-3/2}$$

MAD

Magnetically Arrested Disk

B-flux saturation



$$\dot{M} \sim 4\pi r_B^2 V \rho \sim 5 \times 10^{35} \text{ erg s}^{-1}$$


$$\times \left(\frac{n}{10 \text{ cm}^{-3}} \right) \left(\frac{M}{10 M_{\odot}} \right)^2 \left(\frac{V}{10 \text{ km s}^{-1}} \right)^{-3}$$

$$p_a = \frac{GM\Sigma}{r^2} \sim \frac{GMM\dot{M}}{2\pi r^3 v_r} \Leftrightarrow p_B = \frac{B^2}{8\pi}$$

Bisnovatyi-Kogan & Ruzmaikin 76

Narayan+ 03

$$B_H \sim \sqrt{\frac{4GMM\dot{M}}{r^3 v_r}} \Big|_{r=r_H} \sim 4 \times 10^7 \text{ G} \left(\frac{n}{10 \text{ cm}^{-3}} \right)^{1/2} \left(\frac{V}{10 \text{ km s}^{-1}} \right)^{-3/2}$$



BZ Luminosity

$$L_{BZ} = \frac{\kappa}{4\pi c} \Omega_{BH}^2 \Psi_{BH}^2 \quad \kappa \sim 0.05$$

$$= \frac{\kappa}{4\pi c} \left(\frac{a_* c}{2r_H} \right)^2 \left(\pi r_H^2 B_H \right)^2$$

$$= \frac{\pi \kappa}{16} c a_*^2 r_H^2 B_H^2$$

$$\sim \left(\frac{\kappa}{\varepsilon} \sqrt{\frac{\pi^3}{12} \frac{GM}{c^2 r_H}} \right) a_*^2 \dot{M} c^2$$

$$\sim \dot{M} c^2$$

$$p_B \sim p_a$$

$$B_H \sim \sqrt{\frac{4GM\dot{M}}{r^3 v_r}} \Big|_{r=r_H}$$

$$v_r = \varepsilon v_{ff} = \varepsilon \sqrt{\frac{3GM}{4\pi r}}$$

$$\varepsilon \sim 0.05$$

Bardeen-Petterson Effect

BH spin axis

// Disk axis

“Bardeen-Petterson effect”

// B axis

“Magneto-spin alignment”




Fig. 1. 3D snapshot for an evolved model with $j = 0.99$, initial relative tilt $\theta_{\text{tilt},0} \approx 90^\circ$, and disk thickness $H/R \sim 0.3$. The rotating BH sits at the center of the box of size $r = -40r_g$ to $r = +40r_g$ in each dimension. The snapshot shows the disk near the BH (yellow isosurface, which is mostly flat in the figure plane), the highly magnetized jet region (blue isosurface, with magnetic energy per unit rest-mass energy equal to about 70), the rotational axis of the disk both initially and at large distances (orange cylinder), outer disk (green-yellow volume rendering, more aligned with disk rotational axis at large distances), magnetic field vectors (like iron filings on that surface) for a cross section of the jet (cyan vectors), and jet magnetic field lines (white lines) that trace from the BH out to large distances. The disk and jet near the BH are aligned with the BH spin axis and point mostly in and out of the figure plane, whereas at larger distances the jet points roughly halfway between the BH spin axis and the disk's rotational axis (pointing along the orange cylinder).

Bardeen & Petterson 75
McKinney+ 13

Luminosity Function

$$\frac{dN}{d\dot{M}} = N_{\text{BH}} \int dm_1 \frac{dp(m_1)}{dm_1} \int dm_2 \frac{dp(m_2|m_1)}{dm_2} \int dv \frac{df(v)}{dv} \int dn \frac{d\xi(n)}{dn} \times h(m_1, m_2, v) \delta \left[\dot{M}(n, m_1, m_2, v) - \dot{M} \right],$$


Agol & Kamionkowski 12
KI+ in prep.

BH mass: m_1 : Salpeter, m_2 : Flat, $5M_\odot < m_2 < m_1 < 50M_\odot$

Velocity: Maxwell distribution

+ GW recoil + ISM sound velocity

Density: 5 phases of ISM



Phase	n_1 [cm ⁻³]	n_2 [cm ⁻³]	β	ξ_0	c_s [km s ⁻¹]	H_d
Molecular clouds	10^2	10^5	2.8	10^{-3}	10	75 pc
Cold H _I	10	10^2	3.8	0.04	10	150 pc
Warm H _I	0.3	—	—	0.35	10	0.5 kpc
Warm H _{II}	0.15	—	—	0.2	10	1 kpc
Hot H _{II}	0.002	—	—	0.4	150	3 kpc

Luminosity Function

$$\frac{dN}{d\dot{M}} = N_{\text{BH}} \int dm_1 \frac{dp(m_1)}{dm_1} \int dm_2 \frac{dp(m_2|m_1)}{dm_2} \int dv \frac{df(v)}{dv} \int dn \frac{d\xi(n)}{dn} \\ \times h(m_1, m_2, v) \delta \left[\dot{M}(n, m_1, m_2, v) - \dot{M} \right],$$

Mass function

Agol & Kamionkowski 12

$$\frac{dp(m_1)}{dm_1} = C m_1^{-2.35}, \quad \frac{dp(m_2|m_1)}{dm_2} = \frac{1}{m_1 - M_{\text{min}}}, \quad 5M_{\odot} \leq m_2 \leq m_1 \leq 50M_{\odot}$$

Velocity distribution

$$\frac{df(v)}{dv} = \sqrt{\frac{2}{\pi}} \frac{v^2}{\sigma_v^3} \exp\left(-\frac{v^2}{2\sigma_v^2}\right)$$

$\sigma_v = 40 \text{ km/s}$: isolated binary
 $\sigma_v = 200 \text{ km/s}$: stellar cluster

c_s & v

- $c_s(\text{ISM}) \sim 10 \text{ km s}^{-1}$
- **Formation channel**
 - Massive binary
 - Direct collapse $\Rightarrow$ Little kick
 - SN kick, if $v \propto p$

$$\sigma_{\text{BH}} \sim \frac{M_{\text{NS}}}{M_{\text{BH}}} \sigma_{\text{NS}} \sim \frac{1.4}{60} \times 300 \sim 7 \text{ km s}^{-1}$$
 - 3 body in clusters
 - Escape velocity $\sim 50 \text{ km s}^{-1}$
 - $v \sim v_{\text{Galaxy}} \sim 200 \text{ km s}^{-1}$
- **Obs. of BH candidates**
 - few tens of km s^{-1} or less





Figure 3. Inferred peculiar velocity as a function of black hole mass. Black points denote low-mass X-ray binaries, and the red point represents the high-mass X-ray binary Cygnus X-1. A larger sample is required to make robust inferences about any potential correlation between black hole (or companion) mass and natal kicks.

GW Recoil Velocity

$$v = A \times \eta^2 \sqrt{1 - 4\eta} (1 + B\eta) \cong 62 \text{ km s}^{-1}$$

$$\eta \equiv \frac{q}{(1+q)^2}, \quad q = \frac{M_1}{M_2}$$

$$A = 1.20 \times 10^4 \text{ km s}^{-1}, \quad B = -0.93$$



We assume
 $a_* \sim 0$

Density Distribution

$$\frac{d\xi(n)}{dn} = D\xi_0 n^{-\beta}$$

Volume
filling fraction

Scale
height

Phase	n_1 [cm ⁻³]	n_2 [cm ⁻³]	β	ξ_0	c_s [km s ⁻¹]	H_d
Molecular clouds	10 ²	10 ⁵	2.8	10 ⁻³	10	75 pc
Cold H _I	10	10 ²	3.8	0.04	10	150 pc
Warm H _I	0.3	—	—	0.35	10	0.5 kpc
Warm H _{II}	0.15	—	—	0.2	10	1 kpc
Hot H _{II}	0.002	—	—	0.4	150	3 kpc

$$h(m_1, m_2, v) = \min\left(1, \frac{H_d}{H(v_z)}\right), \quad v_z^2 = \frac{1}{3}(v^2 + v_g^2)$$


$$Z \sim 180 \text{ pc}$$

$$K = 48 M_\odot / \text{pc}^2$$

$$F = 0.01 M_\odot / \text{pc}^3$$

$$\frac{1}{2} v_z^2 = \Phi_z[H(v_z)], \quad \frac{\Phi_z(z)}{2\pi G} = K\left(\sqrt{z^2 + Z^2} - Z\right) + Fz^2$$


Luminosity Function



The most
luminous
BH jet is
 $\sim 10^{36} \text{ erg/s}$
in cold H_I

v_{GW} reduces
 L_j by ~ 10

Particle Acceleration



- Hillas condition

$$E < ZqBR$$

- $L_B \sim 4\pi R^2 (B^2/8\pi) c$
 $\propto (BR)^2$ Blandford 00
 Waxman 04

- $E_{\text{max}} > \text{PeV}$


PeVatron!!!

Barkov+ 12
 KI+ in prep.

Galaxy
 Cluster

Dark
 Matter?

Total Power




BHs $\Leftrightarrow$ Cosmic rays beyond PeV?

$$\frac{\epsilon_{\text{CR}} E_{\text{SN}}}{100\text{yr}} \sim 3 \times 10^{40} \text{ erg s}^{-1}$$

If leptonic $\Leftrightarrow e^{\pm}$ excess?

PAMELA

Positron excess above the predicted secondary



⇒ New sources

– Dark Matter?

– Astrophysical?

⇒ Many papers $> 10^3$



Jul 06 - Feb 08

151672 e^- , 9430 e^+

Astrophysical Models

Pulsar

$$E_{\text{rot}} \sim \frac{1}{2} I \Omega^2 \sim 10^{46} \text{ erg} \left(\frac{P}{\text{sec}} \right)^{-2}$$

Supernova remnant

Microquasar

Gamma-ray burst






White dwarf pulsar

Energy required for $e^\pm$ excess

$$R_{e^+} \sim \frac{\rho_{e^+}}{\rho_p} \frac{t_{\text{esc}}}{t_{\text{cool}}} R_p \sim \frac{10^{46} \text{ erg}}{100 \text{ yr}}$$

TeV Gamma-Ray Sky

HESS 1307.4690



***unIDs dominate
TeV γ -ray sky***

Spatially extended

$$R \sim \theta d \sim 3\text{pc} \left(\frac{\theta}{0.2^\circ} \right) \left(\frac{d}{\text{kpc}} \right)$$

TeV unID

Disk $\Rightarrow$ Galactic origin $d \sim 1-10 \text{ kpc}$

$$F \sim 10^{-11} - 10^{-12} \text{ erg s}^{-1} \text{ cm}^{-2}$$

$$L = 4\pi d^2 F \sim 10^{34-35} \text{ erg s}^{-1} \left(\frac{d}{10 \text{ kpc}} \right)^2$$

$$N \sim 10 - 100$$

Extended

$$R \sim \theta d \sim 30 \text{ pc} \left(\frac{\theta}{0.2^\circ} \right) \left(\frac{d}{10 \text{ kpc}} \right)$$

BH Jet Nebula




Poynting jet
⇒ Dissipation
~ Pulsar wind nebula

Jet termination radius

$$\frac{L}{4\pi R_h^2 \theta^2 c} \sim \rho c_s^2$$

$$R_h \sim 10^{19} \text{ cm} \sim 3 \text{ pc}$$

Log N – Log F



BHs $\Leftrightarrow$ TeV unIDs?

Flux dis. is similar

BH nebula size:

$$\frac{L_j}{4\pi r_h^2 \theta^2 c} \sim \rho V^2$$

$$\Rightarrow r_h \sim 3 \text{ pc}$$

Model Uncertainties

	Number N_{BH}	Velocity σ_v [km s $^{-1}$]	Spin a_*^i	Disk s	Duty cycle $\mathcal{D}$	Total power P_{tot} [erg s $^{-1}$]
Isolated binary (fiducial)	7×10^4	40	0	0	1	$\sim 10^{37}$
Stellar cluster	—	200	—	—	—	$\sim 10^{37} \times 10^{-2}$
High spin	10^8	—	0.2	—	—	$\sim 10^{37} \times 10^3$
Wind	—	—	—	1	—	$\sim 10^{37} \times 10^{-1}$
Feedback	—	—	—	1	10^{-2}	$\sim 10^{37} \times 10^{-3}$

KI, Matsumoto, Teraki, Kashiyaama & Murase in prep.


$$\dot{M}_{BH} \sim \dot{M} \left(\frac{20r_S}{r_{\text{disk}}} \right)^s \quad \text{for } r_{\text{disk}} > 20r_S$$

Blandford & Begelman 99
Yuan+ 16

$$\text{Duty cycle} \sim \frac{t_{\text{active}}}{t_{\text{dormant}}} \sim \frac{\text{Accretion time@}r_{\text{Bondi}}}{\text{Nebula lifetime}} \sim 10^{-2}$$

Uncertainties in total power $\sim \times 10^{\pm 3}$

Dependence on Binary Formation Scenario



$$\dot{M} \propto V^{-3} \sim$$


$$\left(\frac{40 \text{ km s}^{-1}}{200 \text{ km s}^{-1}} \right)^3$$

$$\sim \frac{1}{100}$$

Still relevant
to TeV unIDs

Dependence on Accretion

Disk Model



Adiabatic
inflow-outflow
solution (ADIOS)

$$\dot{M}_{\text{BH}} \approx \dot{M} \left(\frac{20r_{\text{S}}}{r_{\text{disk}}} \right)^s$$

Yuan+ 15

Most ISM does
not accrete to BH

Tip of Iceberg

Gravitational waves

X-ray binary

Cosmic ray?

TeV unID?

Galactic BHs



Summary

- **Merged BHs have huge energy**
- **E_{rot} is extracted by B**
 - ADAF, MAD
 - Blandford-Znajek jet $\Rightarrow$ Luminosity function
- **PeVatron**
- **High energy remnant like PWN**
 - $>\text{PeV}$ CRs
 - TeV unID
 - Positron excess ...


Source Parameters

TABLE I. Source parameters for GW150914. We report median values with 90% credible intervals that include statistical errors, and systematic errors from averaging the results of different waveform models. Masses are given in the source frame; to convert to the detector frame multiply by $(1+z)$ [90]. The source redshift assumes standard cosmology [91].

Primary black hole mass	$36^{+5}_{-4} M_{\odot}$
Secondary black hole mass	$29^{+4}_{-4} M_{\odot}$
Final black hole mass	$62^{+4}_{-4} M_{\odot}$
Final black hole spin	$0.67^{+0.05}_{-0.07}$
Luminosity distance	$410^{+160}_{-180} \text{ Mpc}$
Source redshift z	$0.09^{+0.03}_{-0.04}$

Primary black hole mass	$14.2^{+8.3}_{-3.7} M_{\odot}$
Secondary black hole mass	$7.5^{+2.3}_{-2.3} M_{\odot}$
Chirp mass	$8.9^{+0.3}_{-0.3} M_{\odot}$
Total black hole mass	$21.8^{+5.9}_{-1.7} M_{\odot}$
Final black hole mass	$20.8^{+6.1}_{-1.7} M_{\odot}$
Radiated gravitational-wave energy	$1.0^{+0.1}_{-0.2} M_{\odot} c^2$
Peak luminosity	$3.3^{+0.8}_{-1.6} \times 10^{56} \text{ erg/s}$
Final black hole spin	$0.74^{+0.06}_{-0.06}$
Luminosity distance	$440^{+180}_{-190} \text{ Mpc}$
Source redshift z	$0.09^{+0.03}_{-0.04}$

Event Rate



$9-240 \text{ Gpc}^{-3} \text{yr}^{-1}$

$70 \text{ Gpc}^{-3} \text{yr}^{-1}$

$\div \sim 0.01 \text{ galaxy Mpc}^{-3}$

$\times \sim 10^{10} \text{ yr}$

$\sim 7 \times 10^4$

BHs/galaxy

Black Hole Candidates

BH candidates are discovered as X-ray novae
Half of them have no counterpart

Table 1 Twenty confirmed black holes and twenty black-hole candidates^a

Coordinate Name	Common ^b Name/Prefix	Year ^c	Spec.	P_{orb} (hr)	$f(M)$ ($M_{\odot}$)	M_1 ($M_{\odot}$)
0422+32	(GRO J)	1992/1	M2V	5.1	1.19 ± 0.02	3.7–5.0
0538–641	LMC X–3	–	B3V	40.9	2.3 ± 0.3	5.9–9.2
0540–697	LMC X–1	–	O7III	93.8 ^d	0.13 ± 0.05^d	4.0–10.0: ^e
0620–003	(A)	1975/1 ^f	K4V	7.8	2.72 ± 0.06	8.7–12.9
1009–45	(GRS)	1993/1	K7/M0V	6.8	3.17 ± 0.12	3.6–4.7: ^e
1118+480	(XTE J)	2000/2	K5/M0V	4.1	6.1 ± 0.3	6.5–7.2
1124–684	Nova Mus 91	1991/1	K3/K5V	10.4	3.01 ± 0.15	6.5–8.2
1354–64 ^g	(GS)	1987/2	GIV	61.1 ^g	5.75 ± 0.30	–
1543–475	(4U)	1971/4	A2V	26.8	0.25 ± 0.01	8.4–10.4
1550–564	(XTE J)	1998/5	G8/K8IV	37.0	6.86 ± 0.71	8.4–10.8
1650–500 ^b	(XTE J)	2001/1	K4V	7.7	2.73 ± 0.56	–
1655–40	(GRO J)	1994/3	F3/F5IV	62.9	2.73 ± 0.09	6.0–6.6
1659–487	GX 339–4	1972/10 ⁱ	–	42.1 ^{j,k}	5.8 ± 0.5	–
1705–250	Nova Oph 77	1977/1	K3/7V	12.5	4.86 ± 0.13	5.6–8.3
1819.3–2525	V4641 Sgr	1999/4	B9III	67.6	3.13 ± 0.13	6.8–7.4
1859+226	(XTE J)	1999/1	–	9.2: ^e	$7.4 \pm 1.1:e$	7.6–12.0: ^e
1915+105	(GRS)	1992/Q ^j	K/MIII	804.0	9.5 ± 3.0	10.0–18.0
1956+350	Cyg X–1	–	O9.7Iab	134.4	0.244 ± 0.005	6.8–13.3
2000+251	(GS)	1988/1	K3/K7V	8.3	5.01 ± 0.12	7.1–7.8
2023+338	V404 Cyg	1989/1 ^f	K0III	155.3	6.08 ± 0.06	10.1–13.4

1524–617	(A)	1974/2	–	–	–	–
1630–472	(4U)	1971/15	–	–	–	–
1711.6–3808	(SAX J)	2001/1	–	–	–	–
1716–249	(GRS)	1993/1	–	14.9	–	–
1720–318	(XTE J)	2002/1	–	–	–	–
1730–312	(KS)	1994/1	–	–	–	–
1737–31	(GRS)	1997/1	–	–	–	–
1739–278	(GRS)	1996/1	–	–	–	–
1740.7–2942	(1E)	–	–	–	–	–
1743–322	(H)	1977/4	–	–	–	–
1742–289	(A)	1975/1	–	–	–	–
1746–331	(SLX)	1990/2	–	–	–	–
1748–288	(XTE J)	1998/1	–	–	–	–
1755–324	(XTE J)	1997/1	–	–	–	–
1755–338	(4U)	1971/Q ^j	–	4.5	–	–
1758–258	(GRS)	1990/Q ^j	–	–	–	–


(Continued)

Table 1 Twenty confirmed black holes and twenty black-hole candidates^a

Coordinate Name	Common ^b Name/Prefix	Year ^c	Spec.	P_{orb} (hr)	$f(M)$ ($M_{\odot}$)	M_1 ($M_{\odot}$)
1846–031	(EXO)	1985/1	–	–	–	–
1908+094	(XTE J)	2002/1	–	–	–	–
1957+115	(4U)	–	–	9.3	–	–
2012+381	(XTE J)	1998/1	–	–	–	–

Fermi γ -ray Burst Monitor

GBM detectors at 150914 09:50:45.797 +1.024s



$>50\text{keV}$

0.4s after GW


$T \sim 1\text{sec}$

False alarm ~ 0.0022

$L \sim 1.8^{+1.5}_{-1.0} \text{e}49 \text{erg/s}$

Short GRB!?

No-Go?



$$t_{ff} \sim \frac{1}{\sqrt{G\rho}} \sim 1 \text{ sec}$$

$$\Rightarrow r \sim 10^9 \text{ cm}$$

$$t_m \sim \frac{5}{256} \frac{c^5 r^4}{G^3 M^2 \mu}$$


$$\sim 3 \times 10^3 \text{ sec}$$

$$Lt_m \sim 3 \times 10^{52} \text{ erg}$$

$$> \frac{GMm}{r} \sim 2 \times 10^{52} \text{ erg} \left(\frac{m}{1M_{\text{sun}}} \right)$$

Accretion occurred $\Rightarrow$ Outflow $\Rightarrow$ Unbound

Cooling Function




$$\Lambda \sim 10^{-22} \text{ erg cm}^3 \text{ s}^{-1}$$

$$\Lambda \cdot n^2 \cdot \frac{4\pi}{3} R^3 \sim L$$

$$R \sim 10^{19} \text{ cm} \left(\frac{L}{10^{36} \text{ erg s}^{-1}} \right)^{1/3} n^{-2/3}$$

$$\sim 3 \text{ pc}$$

**ISM can tolerate
the energy injection**



Recoil with Spin

$$v_k = \sqrt{v_m^2 + 2v_m v_\perp \cos \xi + v_\perp^2 + v_{//}^2}$$

$$v_m = A\eta^2 \frac{1-q}{1+q} (1+B\eta)$$

$$v_\perp = H\eta^2 \Delta_{//}$$

$$v_{//} = 16\eta^2 \left[\Delta_\perp \left(V_{11} + 2V_A \tilde{\chi}_{//} + 4V_B \tilde{\chi}_{//}^2 + 8V_C \tilde{\chi}_{//}^3 \right) + 2\tilde{\chi}_\perp \Delta_{//} (C_2 + 2C_3 \tilde{\chi}_{//}) \right] \cos \Theta$$

$$q = \frac{m_2}{m_1}$$

$$M = m_1 + m_2$$

$$\eta = \frac{m_1 m_2}{M^2} = \frac{q}{(1+q)^2}$$

$$S_i = m_i^2 \chi_i$$

$$\Delta = \frac{q\chi_2 \hat{\mathbf{S}}_2 - \chi_1 \hat{\mathbf{S}}_1}{1+q}$$

$$\tilde{\chi} = \frac{q^2 \chi_2 \hat{\mathbf{S}}_2 + \chi_1 \hat{\mathbf{S}}_1}{(1+q)^2}$$

$$\tilde{\chi}_{//} = \tilde{\chi} \cdot \hat{\mathbf{L}}$$

$$\tilde{\chi}_\perp = |\tilde{\chi} \times \hat{\mathbf{L}}|$$

$$A = 1.2 \times 10^4 \text{ km s}^{-1}$$

$$B = -0.93$$

$$H = 6.9 \times 10^3 \text{ km s}^{-1}$$

$$V_{11} = 3677.76 \text{ km s}^{-1}$$

$$V_A = 2481.21 \text{ km s}^{-1}$$

$$V_B = 1792.45 \text{ km s}^{-1}$$

$$V_C = 1506.52 \text{ km s}^{-1}$$

$$C_2 = 1140 \text{ km s}^{-1}$$

$$C_3 = 2481 \text{ km s}^{-1}$$

$$\xi = 145^\circ$$

$$q \sim 1, \chi_2 \sim 0.2$$

$$v_\perp \sim 40 \text{ km/s}$$

$$v_{//} \sim 260 \text{ km/s}$$