

# X-ray Spectrum & Radiative Acceleration of Expanding Pair Fireball in magnetar bursts

ICRR, University of Tokyo,



# TOMOKI WADA



collaborator: Katsuaki Asano

231102マルチメッセンジャー天文学の展開

# Fast Radio Burst

## Brightest radio transient in the universe!

- Short Duration  $\Delta t \sim \mathcal{O}(\text{ms})$
- Radio Band 150 MHz – 8 GHz
- Bright  $L \sim 10^{41} \text{ erg s}^{-1}$
- Cosmological  $D_L \sim 4 \text{ Mpc} - z \sim 2$



(Thornton et al. 2013)

(Petroff et al. 2021)


# FRB 20200428A

FRB & X-ray short burst from  
a galactic magnetar SGR 1935+2154!


FRB luminosity

$$L_{\text{FRB}} \sim 10^{38} \text{ erg s}^{-1}$$

Fainter than others




Spectral luminosity



(Bochenek et al. 2020)

# FRB & X-ray association

count/ms (X-ray)




**FRB !!**

Dim FRB & X-ray short burst from galactic magnetar  
(SGR 1935+2154)  
-> Connection between magnetar burst & FRB

# X-ray short burst associated with FRB

High cut-off energy  $E^{-\alpha} \exp(-E/E_{\text{cut}})$



$$E_{\text{cut}} \sim 80 \text{ keV}$$

c.f., Trapped fireball model

$$T_{\text{eff}} \sim 8 \text{ keV } B^{1/3} R_6^{-1/3} g_{*,14}^{1/6}$$

# High-temperature of X-ray & Radio burst

$$E_{\text{cut}} \sim 80 \text{ keV}$$

Relativistic motion  
of outflow

$$\Gamma \propto r^{3/2}$$

$$T \propto r^{-3/2}$$

- Observed temperature


$$T_{\text{obs}} \sim \Gamma T = T_0$$

↑  
Doppler shift

High  $T_{\text{obs}}$  for high initial  $T_0$


$$E_{\text{FRB}} \sim 10^{-3} E_X$$

Kinetic Energy of outflow  
Converted to radio burst  
@ outer region



## Fireball expanding along flux tube of a magnetar

1. Strong  $\vec{B}$ 
  - number density
  - cross section (E-mode)
2. Baryon loading
3. Lateral diffusion of photons
4. Radiative acceleration w/resonant scattering



# Fireball dynamics

Initially (optically thick),  
Fireball acceleration &  
cooling

$$\Gamma \propto r^{3/2}$$


$$T \propto r^{-3/2}$$

In optically thin region  
Radiative force

$$F_{\text{rad}} \propto \frac{\sigma F}{c}$$

$\sigma$  : cross section

$F$  : X-ray flux






# Normal mode in magnetized pair plasma

There are two modes in magnetized plasma


## X-mode

$$\vec{E}_{\text{wave}} \perp (\vec{k} \times \vec{B}_{\text{bg}})$$



## O-mode

$$\vec{E}_{\text{wave}} \text{ in } \vec{k}, \vec{B}_{\text{bg}} \text{ plane}$$



$$\sigma_E \sim \sigma_T \times \min(1, \omega^2 / \omega_B^2)$$

$$+ \sigma_{\text{res}} \delta(\omega - \omega_B)$$

$$\sigma_O \sim \sigma_T$$


# Radiative acceleration @optically thin region

In optically thin region  
Radiative force

$$F_{\text{rad}} \propto \frac{\sigma_{\text{res}} F}{c}$$

$\sigma_{\text{res}}$  : resonant cross section  
 $F$  : X-ray flux

- ➔ Acceleration by Resonant scattering
- ➔ High kinetic energy



# Basic Equations

Plasma : spherically symmetric fluid equation,  
 particle are ground state of Landau level ( $p_{\perp} = 0$ )

**Solve**

**a steady flow**

~~$$\partial_t (r^2 \rho \Gamma) + \partial_r (r^2 \rho \Gamma \beta) = 0,$$~~

~~$$\partial_t [r^2 (\rho h_{\parallel} \Gamma^2 - p_{\parallel})] + \partial_r [r^2 \rho h_{\parallel} \Gamma^2 \beta] = r^2 G^t,$$~~

~~$$\partial_t [r^2 \rho h_{\parallel} \Gamma^2 \beta] + \partial_r [r^2 (\rho h_{\parallel} \Gamma^2 \beta^2 - p_{\parallel})] = r^2 G^r,$$~~

$$h_{\parallel} = 1 + \frac{e_{\text{th}}}{\rho} + \frac{p_{\parallel}}{\rho}, \quad p_{\parallel} = (\hat{\Gamma} - 1)e_{\text{th}}, \quad \hat{\Gamma} = 3$$

**Radiative  
Force**

**Radiation : Radiation transfer equation,  
 solved by Monte-Carlo scheme**

$$p^{\mu} \left( \frac{\partial}{\partial x^{\mu}} - \Gamma_{\mu\nu}^{\rho} p^{\nu} \frac{\partial}{\partial p^{\rho}} \right) F(x, p) = \left( \frac{dF}{d\tau} \right)_{\text{coll}}$$

$$\left( \frac{dF}{d\tau} \right)_{\text{coll}} = n(x) \left[ -\kappa(x, p) F(x, p) + \int dP' \kappa(x, p') \zeta(x; p' \rightarrow p) F(x, p') \right],$$

# Physical & numerical setup

**Initial condition :**

Analytic solution of fireball

**Optically thin region:**

Solved numerically

Solve radiation in  
a given plasma profile

↓ Radiative force


Solve the plasma

↓ Plasma profile


↑ Radiative force

Solve the radiation

Iteration  
-> steady solution




# Result: radiative acceleration



Acceleration  
by radiation force


$$\Gamma \sim \times 100$$



Energy budget of  
the FRB?

Radiative force acceleration  
-> ultra relativistic outflow

# Result: X-ray spectrum



X-ray is weakly Comptonized

-> Observed spectrum can be created


# Neutrino emission

w/ baryon precursor of fireball

@ outer region



outflow will form shock

-> Particle acceleration



-> neutrino emission

Non-relativistic matter precursor





# Summary

- We study the X-ray spectrum & outflow kinetic luminosity in magnetar short bursts.
- Relativistic outflow is strongly accelerated by the radiation via the resonant scattering.
- X-ray is weakly Comptonized.  
It can be the origin of the observed X-ray spectrum.
- Ultra relativistic outflow can emit neutrinos.

