



# Recent results from atmospheric neutrino analysis at the Super-Kamiokande

**Hiroshi Kaji**

Research Center for Cosmic Neutrinos


Institute for Cosmic Ray Research

University of Tokyo

on behalf of Super-Kamiokande collaboration



Flight length is usually described by zenith angle,  $\theta$ .

$\cos\theta=1$  (down going)




$\cos\theta=-1$  (up going)  
 $L \sim 13000 \text{ km}$

# Atmospheric neutrino



proton



Typically,  $\nu_\mu : \nu_e = 2 : 1$

## Famous as the disappearance of the $\nu_\mu$ flux


- The neutrino oscillation is firstly observed from this sample ( $\theta_{23}, \Delta m_{23}^2$ )
- It sets the best limit to oscillation parameter,  $\theta_{23}$ .

## The sub-dominant oscillation in the $\nu_e$ flux

- Solar term ( $\theta_{12}, \Delta m_{12}^2$ )
- Matter effect term ( $\theta_{13}$ )
- Interference ( $\theta_{12}, \theta_{13}, \Delta m_{12}^2, \delta_{CP}$ )

**Atmospheric- $\nu$  sample is sensitive to all oscillation parameters.**

# Super-Kamiokande



The Super-Kamiokande (SK) is the world's largest water Cherenkov detector.

- Target mass 22.5kton (Fiducial volume)
- Inner detector ~11000 20inch-PMTs
- Outer detector ~2000 8inch-PMTs
- The operation is started since 1996

Run-II : half PMT density

Run-IV (now) : with new electronics

The atmospheric, solar, and Super Nova (relic) neutrinos are studied.

The proton-decay search is still on-going.

The far detector of Japanese LBL  $\nu$  experiments, K2K and T2K.


The atmospheric  $\nu$  data used also for the control sample of both experiments.




# Atmospheric $\nu$ event category

We collect four kinds of events. They have different  $\nu$  energy.


Fully contained (FC)





Partially contained (PC)



Up- $\mu$   
stopping



Up- $\mu$   
through-going



The large energy range is covered with these four kinds of event samples.

# Zenith angle distribution

—  $\nu_\mu - \nu_\tau$  oscillation (best fit)  
 — null oscillation

Live time:

SK-I

1489days (FCPC)

1646days (Up- $\mu$ )

SK-II

799days (FCPC)

827days (Up- $\mu$ )

SK-III

518days (FCPC)


636days (Up- $\mu$ )

The samples with  $E \sim 1$  GeV are further divided to improve sensitivity to low-energy oscillation effects. Totally 420 bins are used in each SK run period.

Momentum


e-like

$\mu$ -like



# Two flavor analysis

$$P(\nu_\alpha \rightarrow \nu_\alpha) = 1 - \sin^2 2\theta \cdot \sin\left(\frac{1.27 \Delta m^2 L}{E}\right)$$



Two analyses are performed at SK.

- Zenith angle analysis
- L/E analysis ← Better sensitivity to  $\Delta m^2$

Allowed Region:

## Zenith angle analysis

$$\Delta m_{23}^2 = 2.11 + 0.11 / -0.19 \times 10^{-3}$$

$$\sin^2 2\theta_{23} > 0.96 \text{ (90\% C.L.)}$$

## L/E analysis

$$\Delta m_{23}^2 = 2.19 + 0.14 / -0.13 \times 10^{-3}$$

$$\sin^2 2\theta_{23} > 0.96 \text{ (90\% C.L.)}$$

**Consistent results!**

**The current best limit to  $\theta_{23}$  is provided.**

# Full three flavor analysis

## “Full three flavor analysis”

takes into account the matter effect, solar, and their interference terms. (Analysis which takes into account only one of matter or solar terms is called “Three flavor analysis”.)

Difference in # of electron events:


$$\Delta_e \equiv \frac{N_e}{N_e^0} \cong \Delta_1(\theta_{13}) \leftarrow \text{Matter} + \Delta_2(\Delta m_{12}^2) \leftarrow \text{Solar} + \Delta_3(\theta_{13}, \Delta m_{12}^2, \delta) \leftarrow \text{Interference}$$

The  $\theta_{13}$  and CP phase,  $\delta_{CP}$ , are studied in this analysis. The interesting behaviors are seen in the  $\nu_e$  flux if sub-dominant terms are considered.

- The  $\theta_{13}$  is in the matter and interference terms.
- The interference term includes  $\delta_{CP}$ .

(The  $\sin^2\theta_{12}$  and  $\Delta m_{12}^2$  are fixed to be 0.304 and  $7.66 \times 10^{-5} \text{ eV}^2$ .)


The resultant  $\theta_{23}$  and  $\Delta m_{23}^2$  are confirmed.  
 $\Rightarrow$  see next slide.



# Confirmation of $\theta_{23}$ and $\Delta m_{23}^2$ results

The resultant  $\theta_{23}$  and  $\Delta m_{23}^2$  are consistent with both SK's two flavor and global three-flavor analyses.

$\chi^2 - \chi^2_{\min}$  distributions



SK Full 3-flavor (NH)

$$(1.88 < \Delta m_{23}^2 < 2.75) \times 10^{-3}$$

$$0.406 < \sin^2 \theta_{23} < 0.629$$

$$(0.93 < \sin^2 2\theta_{23})$$

Global 3-flavor


New J.Phys. 10 113001 (2008)

$$(2.22 < \Delta m_{23}^2 < 2.60) \times 10^{-3}$$

$$0.401 < \sin^2 \theta_{23} < 0.615$$


$$(0.95 < \sin^2 2\theta_{23})$$

Comparison of 3-flavor and 2-flavor




# The $\sin^2\theta_{13}$ and $\delta_{CP}$ results – Normal hierarchy

Normal hierarchy



NH case is more sensitive to  $\sin^2\theta_{13}$  than IH case.

Normal hierarchy





SK has sensitivity to  $\delta_{CP}$  when large  $\theta_{13}$ .

## 1D limit

	Best	90% C.L.
$\sin^2\theta_{13}$	0.006	< 0.066
CP- $\delta$	220°	-

The CHOOZ limit to  $\theta_{13}$  is confirmed.  
No significant constraint to the  $\delta_{CP}$  at 90 % C.L.

# The $\sin^2\theta_{13}$ and $\delta_{CP}$ results – inverted hierarchy



## 1D limit


	Best	90% C.L.
$\sin^2\theta_{13}$	0.044	< 0.122
CP- $\delta$	220°	121.4 - 319.1°

The CHOOZ limit to  $\theta_{13}$  is confirmed.  
 No significant constraint to the  $\delta_{CP}$  at 90 % C.L.


There is a possibility to make the first  $\delta_{CP}$  result with “current” experiments. SK can make a constraint to  $\delta_{CP}$  if large  $\theta_{13}$  is determined by LBL or reactor.

# Mass hierarchy test

The  $\chi^2$  is compared between normal and inverted results.



$E > \sim 1\text{GeV}$  samples tend to favor inverted hierarchy.



**Normal hierarchy (NH):**

$$\chi^2_{\min}/\text{dof} = 469.94/416$$

**Inverted hierarchy (IH):**

$$\chi^2_{\min}/\text{dof} = 468.34/416$$

$$\rightarrow \Delta\chi^2 = 1.6$$

The  $\chi^2$  test is in favor of inverted hierarchy.

**However, no significant difference.**

# Summary of full three-flavor result

**Normal hierarchy** ( $\chi^2_{\min}/\text{dof} = 469.94/416$ )

Parameter	Best point	90% C.L. allowed	68% C.L. allowed
$\Delta m^2_{23} (\times 10^3)$	2.11 eV <sup>2</sup>	1.88 - 2.75 eV <sup>2</sup>	1.99 - 2.54 eV <sup>2</sup>
$\sin^2\theta_{23}$	0.525	0.406 - 0.629	0.441 - 0.597
$\sin^2\theta_{13}$	0.006	< 0.066	< 0.036
CP- $\delta$	220°	-	140.8 - 297.3°

**Inverted hierarchy** ( $\chi^2_{\min}/\text{dof} = 468.34/416$ )

Parameter	Best point	90% C.L. allowed	68% C.L. allowed
$\Delta m^2_{23} (\times 10^3)$	2.51 eV <sup>2</sup>	1.98 - 2.81 eV <sup>2</sup>	2.09 - 2.64 eV <sup>2</sup>
$\sin^2\theta_{23}$	0.575	0.426 - 0.644	0.501 - 0.623
$\sin^2\theta_{13}$	0.044	< 0.122	0.0122 - 0.0850
CP- $\delta$	220°	121.4 - 319.1°	165.6 - 280.4°

# Search for CPT violation

+ data (SK-I,II,III)  
 □ MC  $\nu$   
 ■ MC anti- $\nu$

$$P(\nu_\alpha \rightarrow \nu_\beta) \neq P(\bar{\nu}_\beta \rightarrow \bar{\nu}_\alpha)$$

The CPT violation is suggested by

- LSND-MiniBooNE  $\nu_\mu \rightarrow \nu_e$
- MINOS  $\nu_\mu \rightarrow \nu_\mu$

SK can test MINOS result with same oscillation and same L/E range.


We cannot distinguish  $\nu$  and anti- $\nu$  on an event by event.

We take statistical approach. The  $\cos\theta_{\text{zenith}}$  comparison between samples with different  $\nu$ , anti- $\nu$  compositions.

Two flavor scheme with **four oscillation parameters**:

$$P(\nu_\mu \rightarrow \nu_\mu) = 1 - \sin^2 2\theta \cdot \sin\left(\frac{\Delta m^2 L}{E}\right)$$

$$P(\bar{\nu}_\mu \rightarrow \bar{\nu}_\mu) = 1 - \sin^2 2\bar{\theta} \cdot \sin\left(\frac{\Delta \bar{m}^2 L}{E}\right)$$



# Search for CPT violation

+ data (SK-I,II,III)  
 □ MC  $\nu$   
 ■ MC anti- $\nu$

$$P(\nu_\alpha \rightarrow \nu_\beta) \neq P(\bar{\nu}_\beta \rightarrow \bar{\nu}_\alpha)$$

The CPT violation is suggested by

- LSND-MiniBooNE  $\nu_\mu \rightarrow \nu_e$
- MINOS  $\nu_\mu \rightarrow \nu_\mu$

SK can test MINOS result with same oscillation and same L/E range.


We cannot distinguish  $\nu$  and anti- $\nu$  on an event by event.

We take statistical approach. The  $\cos\theta_{\text{zenith}}$  comparison between samples with different  $\nu$ , anti- $\nu$  compositions.

Two flavor scheme with **four oscillation parameters**:

$$P(\nu_\mu \rightarrow \nu_\mu) = 1 - \sin^2 2\theta \cdot \sin\left(\frac{\Delta m^2 L}{E}\right)$$

$$P(\bar{\nu}_\mu \rightarrow \bar{\nu}_\mu) = 1 - \sin^2 2\bar{\theta} \cdot \sin\left(\frac{\Delta \bar{m}^2 L}{E}\right)$$



**Composition differences**


**⇒ Determine  $\nu$  and anti- $\nu$  oscillations, separately.**

# Arrowed region for anti-neutrino

Parameter	Best Fit	90% C.L. Bound
$\Delta m^2$	$2.1 \times 10^{-3} \text{eV}^2$	$[1.8, 2.7] \times 10^{-3} \text{eV}^2$
$\Delta \bar{m}^2$	$2.0 \times 10^{-3} \text{eV}^2$	$[1.5, 3.1] \times 10^{-3} \text{eV}^2$
$\sin^2 2\theta$	1.0	[0.92, 1.0]
$\sin^2 2\bar{\theta}$	1.0	[0.88, 1.0]

The oscillation parameters for  $\nu$  and anti- $\nu$  are consistent. No evidence for CPT violation is observed.

However, there is no inconsistency between the MINOS and our results since partially same region is allowed by both experiments.




# Tau appearance analysis

SK-I only

PRL97,171801 (2006)

( $\sin^2 2\theta = 1.0$ ,  $\Delta m^2 = 2.4 \times 10^{-3} \text{eV}^2$  assumed for MC)

Best fit  $\cos\theta_{\text{zenith}}$  distribution  
after standard+likelihood



	Data	Signal	Background
Fiducial Vol.	-	78.4 (100%)	17135 (100%)
$E_{\text{vis}} > 1330 \text{MeV}$	2888	51.5 (65.7%)	2943 (17.2%)
1st-ring e-like	1803	47.1 (60.1%)	1765 (10.3%)
<b>Likelihood</b>	<b>649</b>	<b>33.8 (43.1%)</b>	<b>647 (3.79%)</b>
<b>Neural Network</b>	<b>603</b>	<b>30.6 (39.0%)</b>	<b>577 (3.36%)</b>

Two approaches are performed.

**Likelihood** ( $\chi^2/\text{dof} = 7.6/8$ )

Best fit:  $N_\tau = 138 \pm 48(\text{stat.})_{-32}^{+15}(\text{syst})$

Expected:  $N_\tau = 78 \pm 26(\text{syst})$

**The  $\tau$  enrich sample is consistent with  $\nu_\mu - \nu_\tau$  oscillation.**  
**The  $2.4 \sigma$  signal is determined.**

The analysis will be updated with SK-I,II,III, soon.

# Summary of results

## Neutrino oscillation

	Data-set	Remarks
Two flavor	SK-I,II,III	Zenith, L/E
Three flavor	SK-I,II,III	Solar term( $\theta_{12}$ , $\Delta m_{12}^2$ ), matter effect( $\theta_{13}$ )
Full three flavor	SK-I,II,III	$\theta_{13}$ , $\delta_{CP}$
CPT violation	SK-I,II,III	No evidence
Non-standard interactions	SK-I,II	No evidence
Tau appearance	SK-I	2.4 $\sigma$ , will be updated soon

## Proton decay



	Data-set	Remarks
$p \rightarrow e^+ \pi^0$	SK-I,II,III	Life time $> 1.0 \times 10^{34}$ years

## Astro-physics



	Data-set	Remarks
WIMP search	SK-I,II,III	Galactic Center, Diffuse source

# SK-IV preliminary result

e-like




$\mu$ -like



— SK-IV 449 days data

—  $\nu_\mu - \nu_\tau$  oscillation

— null oscillation



SK-IV operation  
is stable and  
successful.

- This result is released for supporting the T2K analysis.
- The oscillation analysis is on-going and will be released in next summer.

# Conclusion

Super-Kamiokande plays important roles for the  $\nu$ -oscillation analysis.

## **Atmospheric- $\nu$ sample is sensitive to all oscillation parameters.**

- The best limit to  $\theta_{23}$  is produced.
- We focus on the sub-dominant effects. The CHOOZ limit to  $\theta_{13}$  is confirmed. The CP phase,  $\delta_{CP}$ , analysis is performed.

## **Search for CP violation in $\nu$ -oscillation is no longer future plan.**

- There is a possibility to make the first result by global analysis of “current” experiments.
- The constraint to  $\delta_{CP}$  can be made by atmospheric- $\nu$  sample if large  $\theta_{13}$  is determined by LBL or reactor experiments.

## **There is no evidence for CPT violation at atmospheric- $\nu$ flux.**

- SK tests the “MINOS-type” CPT violation.
- The SK allowed region suggest CPT conservation. However, there is no inconsistency since MINOS also allows this region.